

## Generalized Degree Distance of Strong Product of Graphs

K. Pattabiraman\*, P. Kandan

Department of Mathematics, Annamalai University, Annamalainagar 608 002,  
India.

E-mail: pramank@gmail.com

E-mail: kandan2k@gmail.com

ABSTRACT. In this paper, the exact formulae for the generalized degree distance, degree distance and reciprocal degree distance of strong product of a connected and the complete multipartite graph with partite sets of sizes  $m_0, m_1, \dots, m_{r-1}$  are obtained. Using the results obtained here, the formulae for the degree distance and reciprocal degree distance of the closed and open fence graphs are computed.

**Keywords:** Generalized degree distance, Degree distance, Reciprocal degree distance, Strong product.


**2000 Mathematics subject classification:** 05C12, 05C76.

### 1. INTRODUCTION

All the graphs considered in this paper are simple and connected. For vertices  $u, v \in V(G)$ , the distance between  $u$  and  $v$  in  $G$ , denoted by  $d_G(u, v)$ , is the length of a shortest  $(u, v)$ -path in  $G$  and let  $d_G(v)$  be the degree of a vertex  $v \in V(G)$ . The *strong product* of graphs  $G$  and  $H$ , denoted by  $G \boxtimes H$ , is the graph with vertex set  $V(G) \times V(H) = \{(u, v) : u \in V(G), v \in V(H)\}$  and  $(u, x)(v, y)$  is an edge whenever (i)  $u = v$  and  $xy \in E(H)$ , or (ii)  $uv \in E(G)$  and  $x = y$ , or (iii)  $uv \in E(G)$  and  $xy \in E(H)$ , see Fig.1.

---

\*Corresponding Author


 $C_3 \boxtimes P_3$ 
Fig. 1. Strong product of  $C_3$  and  $P_3$ .

A *topological index* of a graph is a real number related to the graph; it does not depend on labeling or pictorial representation of a graph. In theoretical chemistry, molecular structure descriptors (also called topological indices) are used for modelling physicochemical, pharmacologic, toxicologic, biological and other properties of chemical compounds [8]. There exist several types of such indices, especially those based on vertex and edge distances. One of the most intensively studied topological indices is the Wiener index.

Let  $G$  be a connected graph. Then *Wiener index* of  $G$  is defined as  $W(G) = \frac{1}{2} \sum_{u,v \in V(G)}$   $d_G(u,v)$  with the summation going over all pairs of distinct vertices of  $G$ . This definition can be further generalized in the following way:

$W_\lambda(G) = \frac{1}{2} \sum_{u,v \in V(G)}$   $d_G^\lambda(u,v)$ , where  $d_G^\lambda(u,v) = (d_G(u,v))^\lambda$  and  $\lambda$  is a real

number [9, 10]. If  $\lambda = -1$ , then  $W_{-1}(G) = H(G)$ , where  $H(G)$  is Harary index of  $G$ . In the chemical literature also  $W_{\frac{1}{2}}$  [27] as well as the general case  $W_\lambda$  were examined [6, 11]. Wiener index of 2-dimensional square and comb lattices with open ends is obtained by Graovac et al. in [5]. In [17] the Wiener index of HAC5C7[p, q] and HAC5C6C7[p, q] nanotubes are computed by using GAP program. Dobrynin and Kochetova [4] and Gutman [7] independently proposed a vertex-degree-weighted version of Wiener index called *degree distance* or *Schultz molecular topological index*, which is defined for a connected graph  $G$  as

$DD(G) = \frac{1}{2} \sum_{u,v \in V(G)}$   $(d_G(u) + d_G(v))d_G(u,v)$ , where  $d_G(u)$  is the degree of the

vertex  $u$  in  $G$ . Note that the degree distance is a degree-weight version of the Wiener index. In the literature, many results on the degree distance  $DD(G)$  have been put forward in past decades and they mainly deal with extreme properties of  $DD(G)$ . Tomescu[24] showed that the star is the unique graph with minimum degree distance within the class on  $n$ -vertex connected graphs. Tomescu[25] deduced properties of graphs with minimum degree distance in

the class of  $n$ -vertex connected graphs with  $m \geq n - 1$  edges. For other related results along this line, see [2, 14, 18].

Additively weighted Harary index ( $H_A$ ) or reciprocal degree distance ( $RDD$ ) is defined in [1] as  $H_A(G) = RDD(G) = \frac{1}{2} \sum_{u,v \in V(G)} \frac{(d_G(u) + d_G(v))}{d_G(u,v)}$ . In [12],

Hamzeh et. al recently introduced generalized degree distance of graphs. Hua and Zhang [15] have obtained lower and upper bounds for the reciprocal degree distance of graph in terms of other graph invariants including the degree distance, Harary index, the first Zagreb index, the first Zagreb coindex, pendent vertices, independence number, chromatic number and vertex-, and edge-connectivity. Pattabiraman and Vijayaragavan [21, 22] have obtained the reciprocal degree distance of join, tensor product, strong product and wreath product of two connected graphs in terms of other graph invariants. The chemical applications and mathematical properties of the reciprocal degree distance are well studied in [1, 19, 23].

The generalized degree distance, denoted by  $H_\lambda(G)$ , is defined as

$$H_\lambda(G) = \frac{1}{2} \sum_{u,v \in V(G)} (d_G(u) + d_G(v)) d_G^\lambda(u,v), \text{ where } \lambda \text{ is a any real number.}$$

If  $\lambda = 1$ , then  $H_\lambda(G) = DD(G)$  and if  $\lambda = -1$ , then  $H_\lambda(G) = RDD(G)$ . The generalized degree distance of unicyclic and bicyclic graphs are studied by Hamzeh et. al [12, 13]. Also they are given the generalized degree distance of Cartesian product, join, symmetric difference, composition and disjunction of two graphs. It is well known that many graphs arise from simpler graphs via various graph operations. Hence it is important to understand how certain invariants of such product graphs are related to the corresponding invariants of the original graphs. In this paper, the exact formulae for the generalized degree distance, degree distance and reciprocal degree distance of strong product  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , where  $K_{m_0, m_1, \dots, m_{r-1}}$  is the complete multipartite graph with partite sets of sizes  $m_0, m_1, \dots, m_{r-1}$  are obtained.

The *first Zagreb index* is defined as  $M_1(G) = \sum_{u \in V(G)} d_G(u)^2$ . In fact, one can rewrite the first Zagreb index as  $M_1(G) = \sum_{uv \in E(G)} (d_G(u) + d_G(v))$ . The Zagreb indices are found to have applications in QSPR and QSAR studies as well, see [3].

If  $m_0 = m_1 = \dots = m_{r-1} = s$  in  $K_{m_0, m_1, \dots, m_{r-1}}$  (the complete multipartite graph with partite sets of sizes  $m_0, m_1, \dots, m_{r-1}$ ), then we denote it by  $K_{r(s)}$ . For  $S \subseteq V(G)$ ,  $\langle S \rangle$  denotes the subgraph of  $G$  induced by  $S$ . For two subsets  $S, T \subset V(G)$ , not necessarily disjoint, by  $d_G(S, T)$ , we mean the sum of the distances in  $G$  from each vertex of  $S$  to every vertex of  $T$ , that is,  $d_G(S, T) =$

$$\sum_{s \in S, t \in T} d_G(s, t).$$

2. GENERALIZED DEGREE DISTANCE OF STRONG PRODUCT OF GRAPHS

In this section, we obtain the generalized degree distance of  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ . Let  $G$  be a simple connected graph with  $V(G) = \{v_0, v_1, \dots, v_{n-1}\}$  and let  $K_{m_0, m_1, \dots, m_{r-1}}$ ,  $r \geq 2$ , be the complete multipartite graph with partite sets  $V_0, V_1, \dots, V_{r-1}$  and let  $|V_i| = m_i$ ,  $0 \leq i \leq r-1$ . In the graph  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , let  $B_{ij} = v_i \times V_j$ ,  $v_i \in V(G)$  and  $0 \leq j \leq r-1$ . For our convenience, as in the case of tensor product, the vertex set of  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$  is written as

$$V(G) \times V(K_{m_0, m_1, \dots, m_{r-1}}) = \bigcup_{j=0}^{r-1} B_{ij}.$$

As in the tensor product of graphs, let

$$\mathcal{B} = \{B_{ij}\}_{i=0,1,\dots,n-1, j=0,1,\dots,r-1}.$$

Let  $X_i = \bigcup_{j=0}^{r-1} B_{ij}$  and  $Y_j = \bigcup_{i=0}^{n-1} B_{ij}$ ; we call  $X_i$  and  $Y_j$

as *layer* and *column* of  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , respectively, see Figures 2 and 3. If we denote  $V(B_{ij}) = \{x_{i1}, x_{i2}, \dots, x_{im_j}\}$  and  $V(B_{kp}) = \{x_{k1}, x_{k2}, \dots, x_{km_p}\}$ , then  $x_{i\ell}$  and  $x_{k\ell}$ ,  $1 \leq \ell \leq j$ , are called the *corresponding vertices* of  $B_{ij}$  and  $B_{kp}$ . Further, if  $v_i v_k \in E(G)$ , then the induced subgraph  $\langle B_{ij} \cup B_{kp} \rangle$  of  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$  is isomorphic to  $K_{|V_j||V_p|}$  or,  $m_p$  independent edges joining the corresponding vertices of  $B_{ij}$  and  $B_{kj}$  according as  $j \neq p$  or  $j = p$ , respectively.

Structure of shortest paths in  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$  corresponding to an edge in  $G$ .




Fig. 2.

If  $v_i v_k \in E(G)$ , then shortest paths of length 1 and 2 from  $B_{ij}$  to  $B_{kj}$  are shown in solid edges, where the vertical line between  $B_{ij}$  and  $B_{kj}$  denotes the edge joining the corresponding vertices of  $B_{ij}$  and  $B_{kj}$ . The broken edges denote a shortest path of length 2 from a vertex of  $B_{ij}$  to a vertex of  $B_{ij}$ .

The proof of the following lemma follows easily from the properties and structure of  $G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , see Figs. 2 and 3.

**Lemma 2.1.** Let  $G$  be a connected graph and let  $B_{ij}, B_{kp} \in \mathcal{B}$  of the graph  $G' = G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , where  $r \geq 2$ .

(i) If  $v_i v_k \in E(G)$  and  $x_{it} \in B_{ij}, x_{kt} \in B_{kj}$ , then

$$d_{G'}(x_{it}, x_{kt}) = \begin{cases} 1, & \text{if } t = \ell, \\ 2, & \text{if } t \neq \ell, \end{cases}$$

and if  $x_{it} \in B_{ij}, x_{kt} \in B_{kp}, j \neq p$ , then  $d_{G'}(x_{it}, x_{kt}) = 1$ .

(ii) If  $v_i v_k \notin E(G)$ , then for any two vertices  $x_{it} \in B_{ij}, x_{kt} \in B_{kp}, d_{G'}(x_{it}, x_{kt}) = d_G(v_i, v_k)$ .

(iii) For any two distinct vertices in  $B_{ij}$ , their distance is 2.

The proof of the following lemma follows easily from Lemma 2.1. The lemma is used in the proof of the main theorems of this section.

**Lemma 2.2.** Let  $G$  be a connected graph and let  $B_{ij}, B_{kp} \in \mathcal{B}$  of the graph  $G' = G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ , where  $r \geq 2$ .

(i) If  $v_i v_k \in E(G)$ , then

$$d_{G'}^\lambda(B_{ij}, B_{kp}) = \begin{cases} m_j m_p, & \text{if } j \neq p, \\ (1 - 2^\lambda(m_j - 1))m_j, & \text{if } j = p. \end{cases}$$

(ii) If  $v_i v_k \notin E(G)$ , then  $d_{G'}^\lambda(B_{ij}, B_{kp}) = \begin{cases} m_j m_p d_G^\lambda(v_i, v_k), & \text{if } j \neq p, \\ m_j^2 d_G^\lambda(v_i, v_k), & \text{if } j = p. \end{cases}$

(iii)  $d_{G'}^\lambda(B_{ij}, B_{ip}) = \begin{cases} m_j m_p, & \text{if } j \neq p, \\ 2^\lambda m_j(m_j - 1), & \text{if } j = p. \end{cases}$




Fig. 3.

Corresponding to a shortest path of length  $k > 1$  in  $G$ , the shortest path from any vertex of  $B_{ij}$  to any vertex of  $B_{kj}$  (resp. any vertex of  $B_{ij}$  to any

vertex of

$B_{kp}, p \neq j$ ) of length  $k$  is shown in solid edges (resp. broken edges).

**Lemma 2.3.** *Let  $G$  be a connected graph and let  $B_{ij}$  in  $G' = G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ . Then the degree of a vertex  $(v_i, u_j) \in B_{ij}$  in  $G'$  is  $d_{G'}((v_i, u_j)) = d_G(v_i) + (n_0 - m_j) + d_G(v_i)(n_0 - m_j)$ , where  $n_0 = \sum_{j=0}^{r-1} m_j$ .*

*Remark 2.4.* The sums  $\sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} m_j m_p = 2q, \sum_{j=0}^{r-1} m_j^2 = n_0^2 - 2q, \sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} m_j^2 m_p = n_0^3 - 2n_0q - \sum_{j=0}^{r-1} m_j^3 = \sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} m_j m_p^2$  and  $\sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} m_j^3 m_p = n_0 \sum_{j=0}^{r-1} m_j^3 - \sum_{j=0}^{r-1} m_j^4 = \sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} m_j m_p^3$ , where  $n_0 = \sum_{j=0}^{r-1} m_j$  and  $q$  is the number of edges of  $K_{m_0, m_1, \dots, m_{r-1}}$ .

**Theorem 2.5.** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then  $H_\lambda(G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}) = (n_0^2 + 2n_0q)H_\lambda(G) + 4n_0qW_\lambda(G) + M_1(G)(1 - 2^\lambda) \left( 2qn_0 - n_0^3 - n_0^2 + n_0 + 4q + \sum_{j=0}^{r-1} m_j^3 \right) + m \left( 4qn_0(3 - 2^{\lambda+1}) - 2n_0^3(2 - 2^{\lambda+1}) + 4q(2 - 2^\lambda - 2^{\lambda+1}) + n_0(n_0 - 1)2^{\lambda+1} + 2(2 - 2^{\lambda+1}) \sum_{j=0}^{r-1} m_j^3 \right) + n \left( 2n_0q(2 - 2^\lambda) + n_0^3(2^\lambda - 1) - 2^{\lambda+1}q + (1 - 2^\lambda) \sum_{j=0}^{r-1} m_j^3 \right), r \geq 2$ .*

*Proof.* Let  $G' = G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}$ . Clearly,

$$\begin{aligned}
 H_\lambda(G') &= \frac{1}{2} \sum_{B_{ij}, B_{kp} \in \mathcal{B}} \left( d_{G'}(B_{ij}) + d_{G'}(B_{kp}) \right) d_{G'}^\lambda(B_{ij}, B_{kp}) \\
 &= \frac{1}{2} \left( \sum_{i=0}^{n-1} \sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} \left( d_{G'}(B_{ij}) + d_{G'}(B_{ip}) \right) d_{G'}^\lambda(B_{ij}, B_{ip}) \right. \\
 &\quad + \sum_{\substack{i,k=0 \\ i \neq k}}^{n-1} \sum_{j=0}^{r-1} \left( d_{G'}(B_{ij}) + d_{G'}(B_{kj}) \right) d_{G'}^\lambda(B_{ij}, B_{kj}) \\
 &\quad + \sum_{\substack{i,k=0 \\ i \neq k}}^{n-1} \sum_{\substack{j,p=0 \\ j \neq p}}^{r-1} \left( d_{G'}(B_{ij}) + d_{G'}(B_{kp}) \right) d_{G'}^\lambda(B_{ij}, B_{kp}) \\
 &\quad \left. + \sum_{i=0}^{n-1} \sum_{j=0}^{r-1} \left( d_{G'}(B_{ij}) + d_{G'}(B_{ij}) \right) d_{G'}^\lambda(B_{ij}, B_{ij}) \right) \\
 &= \frac{1}{2} \{A_1 + A_2 + A_3 + A_4\}, \tag{2.1}
 \end{aligned}$$

where  $A_1, A_2, A_3$  and  $A_4$  are the sums of the terms of the above expression, in order.

We shall obtain  $A_1$  to  $A_4$  of (2.1), separately.

$$\begin{aligned}
 A_1 &= \sum_{i=0}^{n-1} \sum_{\substack{j, p=0 \\ j \neq p}}^{r-1} (d_{G'}(B_{ij}) + d_{G'}(B_{ip})) d_{G'}^\lambda(B_{ij}, B_{ip}) \\
 &= \sum_{i=0}^{n-1} \sum_{\substack{j, p=0 \\ j \neq p}}^{r-1} (2d_G(v_i) + d_G(v_i)(2n_0 - m_j - m_p) + (2n_0 - m_j - m_p)) m_j m_p, \\
 &\quad \text{by Lemmas 2.2 and 2.3} \\
 &= 8mq + 2m \left( 4n_0q - 2(n_0^3 - 2n_0q - \sum_{j=0}^{r-1} m_j^3) \right) + n \left( 4n_0q - 2(n_0^3 - 2n_0q - \sum_{j=0}^{r-1} m_j^3) \right), \\
 &\quad \text{by Remark 2.4} \\
 &= 2m \left( 4q + 8n_0q - 2n_0^3 + 2 \sum_{j=0}^{r-1} m_j^3 \right) + n \left( 8n_0q - 2n_0^3 + 2 \sum_{j=0}^{r-1} m_j^3 \right).
 \end{aligned}$$

$$\begin{aligned}
 A_2 &= \sum_{j=0}^{r-1} \sum_{\substack{i, k=0 \\ i \neq k}}^{n-1} (d_{G'}(B_{ij}) + d_{G'}(B_{kj})) d_{G'}^\lambda(B_{ij}, B_{kj}) \\
 &= \sum_{j=0}^{r-1} \sum_{\substack{i, k=0 \\ i \neq k \\ v_i v_k \in E(G)}}^{n-1} ((d_G(v_i) + d_G(v_k)) + 2(n_0 - m_j) + (n_0 - m_j)(d_G(v_i) + d_G(v_k))) \\
 &\quad \times (1 - 2^\lambda + 2^\lambda m_j) m_j \\
 &\quad + \sum_{j=0}^{r-1} \sum_{\substack{i, k=0 \\ i \neq k \\ v_i v_k \notin E(G)}}^{n-1} ((d_G(v_i) + d_G(v_k)) + 2(n_0 - m_j) + (n_0 - m_j)(d_G(v_i) + d_G(v_k))) \\
 &\quad \times m_j^2 d_G^\lambda(v_i, v_k), \\
 &= \sum_{j=0}^{r-1} \sum_{\substack{i, k=0 \\ i \neq k \\ v_i v_k \in E(G)}}^{n-1} ((d_G(v_i) + d_G(v_k)) + 2(n_0 - m_j) + (n_0 - m_j)(d_G(v_i) + d_G(v_k))) \\
 &\quad \times ((1 - 2^\lambda) m_j + (2^\lambda - 1) m_j^2) \\
 &\quad + \sum_{j=0}^{r-1} \sum_{\substack{i, k=0 \\ i \neq k}}^{n-1} ((d_G(v_i) + d_G(v_k)) + 2(n_0 - m_j) + (n_0 - m_j)(d_G(v_i) + d_G(v_k))) \\
 &\quad \times m_j^2 d_G^\lambda(v_i, v_k) \\
 &= 2H_\lambda(G) \left( n_0^3 + n_0^2 - 2q - 2n_0q - \sum_{j=0}^{r-1} m_j^3 \right) + 4W_\lambda(G) \left( n_0^3 - 2n_0q - \sum_{j=0}^{r-1} m_j^3 \right) \\
 &\quad + 2M_1(G) (1 - 2^\lambda) \left( 2qn_0 - n_0^3 - n_0^2 + n_0 + 4q + \sum_{j=0}^{r-1} m_j^3 \right) \\
 &\quad + 4m(1 - 2^\lambda) \left( 2qn_0 - n_0^3 + 2q + \sum_{j=0}^{r-1} m_j^3 \right),
 \end{aligned}$$

by Remark 2.4.

$$\begin{aligned} A_3 &= \sum_{\substack{i, k=0 \\ i \neq k}}^{n-1} \sum_{\substack{j, p=0 \\ j \neq p}}^{r-1} (d_{G'}(B_{ij}) + d_{G'}(B_{kp})) d_{G'}^\lambda(B_{ij}, B_{kp}) \\ &= \sum_{\substack{i, k=0 \\ i \neq k}}^{n-1} \sum_{\substack{j, p=0 \\ j \neq p}}^{r-1} (d_G(v_i) + (n_0 - m_j) + d_G(v_i)(n_0 - m_j) + d_G(v_k) + (n_0 - m_p) \\ &\quad + d_G(v_k)(n_0 - m_p)) m_j m_p d_G^\lambda(v_i, v_k), \end{aligned}$$

by Lemmas 2.2 and Lemma 2.3

$$\begin{aligned} &= \sum_{\substack{i, k=0 \\ i \neq k}}^{n-1} \sum_{\substack{j, p=0 \\ j \neq p}}^{r-1} \left( (d_G(v_i) + d_G(v_k)) d_G^\lambda(v_i, v_k) m_j m_p + d_G(v_i) d_G^\lambda(v_i, v_k) (n_0 - m_j) m_j m_p \right. \\ &\quad \left. + (2n_0 - m_j - m_p) m_j m_p d_G^\lambda(v_i, v_k) + d_G(v_k) d_G^\lambda(v_i, v_k) (n_0 - m_p) m_j m_p \right) \\ &= 2H_\lambda(G) \left( 2q + 4n_0q - n_0^3 + \sum_{j=0}^{r-1} m_j^3 \right) + 2W_\lambda(G) \left( 8n_0q - 2n_0^3 + 2 \sum_{j=0}^{r-1} m_j^3 \right), \end{aligned}$$

by Remark 2.4.

$$\begin{aligned} A_4 &= \sum_{i=0}^{n-1} \left( \sum_{j=0}^{r-1} (d_{G'}(B_{ij}) + d_{G'}(B_{ij})) d_{G'}^\lambda(B_{ij}, B_{ij}) \right) \\ &= \sum_{i=0}^{n-1} \sum_{j=0}^{r-1} 2^{\lambda+1} (d_G(v_i) + (n_0 - m_j) + d_G(v_i)(n_0 - m_j)) m_j (m_j - 1), \text{ by Lemmas 2.2 and 2.3} \\ &= 2^{\lambda+1} \sum_{i=0}^{n-1} \sum_{j=0}^{r-1} (d_G(v_i)(m_j^2 - m_j) + (n_0 - m_j)(m_j^2 - m_j) + d_G(v_i)(n_0 - m_j)(m_j^2 - m_j)) \\ &= 2^{\lambda+1} \left( 2m(n_0^2 - 2q - n_0) + n(n_0^3 - 2qn_0 - n_0^2 - \sum_{j=0}^{r-1} m_j^3 + n_0^2 - 2q) \right. \\ &\quad \left. + 2m(n_0^3 - 2qn_0 - n_0^2 - \sum_{j=0}^{r-1} m_j^3 + n_0^2 - 2q) \right), \text{ by Remark 2.4} \\ &= 2^{\lambda+2} m \left( n_0^3 + n_0^2 - 4q - n_0 - 2qn_0 - \sum_{j=0}^{r-1} m_j^3 \right) + 2^{\lambda+1} n \left( n_0^3 - 2qn_0 - 2q - \sum_{j=0}^{r-1} m_j^3 \right). \end{aligned}$$

Using (2.2), (2), (2.2) and (2.2) in (2.1), we have

$$\begin{aligned} H_\lambda(G') &= (n_0^2 + 2n_0q)H_\lambda(G) + 4n_0qW_\lambda(G) \\ &\quad + M_1(G)(1 - 2^\lambda) \left( 2qn_0 - n_0^3 - n_0^2 + n_0 + 4q + \sum_{j=0}^{r-1} m_j^3 \right) \\ &\quad + m \left( 4qn_0(3 - 2^{\lambda+1}) - 2n_0^3(2 - 2^{\lambda+1}) + 4q(2 - 2^\lambda - 2^{\lambda+1}) + n_0(n_0 - 1)2^{\lambda+1} \right. \\ &\quad \left. + 2(2 - 2^{\lambda+1}) \sum_{j=0}^{r-1} m_j^3 \right) + n \left( 2n_0q(2 - 2^\lambda) + n_0^3(2^\lambda - 1) - 2^{\lambda+1}q + (1 - 2^\lambda) \sum_{j=0}^{r-1} m_j^3 \right). \end{aligned}$$

□

Using  $\lambda = 1$  in Theorem 2.5, we have the following corollary, which is the degree distance of the strong product of graphs.



**Corollary 2.6.** *Let  $G$  be a connected graph with  $n$  vertices. Then  $DD(G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}) = (n_0^2 + 2n_0q)DD(G) + 4n_0qW(G) + M_1(G)(n_0^3 + n_0^2 - 2qn_0 - n_0 - 4q - \sum_{j=0}^{r-1} m_j^3) + 4m(n_0^3 - 4q - n_0q + n_0^2 - n_0 - \sum_{j=0}^{r-1} m_j^3) + n(n_0^3 - 4q - \sum_{j=0}^{r-1} m_j^3)$ ,  $r \geq 2$ .*

If  $m_i = s$ ,  $0 \leq i \leq r - 1$ , in Corollary 2.6, we have the following

**Corollary 2.7.** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then  $DD(G \boxtimes K_{r(s)}) = r^2s^2(rs - s + 1)DD(G) + 2r^2s^3(r - 1)W(G) + M_1(G)rs(rs^2 - rs + 2s - s^2 - 1) + 2mrs(r^2s^2 - 2rs + rs^2 + 4s - 2s^2 - 2) + nrs^2(r^2s - 2r - s + 2)$ ,  $r \geq 2$ .*

As  $K_r = K_{r(1)}$ , the above corollary gives the following

**Corollary 2.8.** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then  $DD(G \boxtimes K_r) = r^3DD(G) + 2r^2(r - 1)W(G) + 2rm(r - 1)^2 + rn(r - 1)^2$ ,  $r \geq 2$ .*

Using  $\lambda = -1$  in Theorem 2.5, we obtain the reciprocal degree distance of strong product of graphs.

**Corollary 2.9.** *Let  $G$  be a connected graph with  $n$  vertices. Then  $RDD(G \boxtimes K_{m_0, m_1, \dots, m_{r-1}}) = (n_0^2 + 2n_0q)RDD(G) + 4n_0qH(G) + \frac{M_1(G)}{2}(n_0(n_0 + 1) - (n_0 + 2)(n_0^2 - 2q) + \sum_{j=0}^{r-1} m_j^3) + m(8n_0q - 2n_0^3 + n_0^2 - n_0 + 2q + 2\sum_{j=0}^{r-1} m_j^3) + \frac{n}{2}(6n_0q - n_0^3 - 2q + \sum_{j=0}^{r-1} m_j^3)$ ,  $r \geq 2$ .*

If  $m_i = s$ ,  $0 \leq i \leq r - 1$ , in Corollary 2.9, we have the following

**Corollary 2.10.** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then  $RDD(G \boxtimes K_{r(s)}) = r^2s^2(rs - s + 1)RDD(G) + 2r^2s^3(r - 1)H(G) + \frac{M_1(G)rs}{2}(rs - rs^2 - 2s + s^2 + 1) + mrs(2r^2s^2 - 4rs^2 + 2rs + 2s^2 - s - 1) + \frac{nr s^2}{2}(2r^2s - 3rs + s - r + 1)$ .*

As  $K_r = K_{r(1)}$ , the above corollary gives the following

**Corollary 2.11.** *Let  $G$  be a connected graph with  $n$  vertices and  $m$  edges. Then  $RDD(G \boxtimes K_r) = r(r^2RDD(G) + 2r(r - 1)H(G) + 2r(r - 1)m + n(r - 1)^2)$ .*

As an application we present formulae for degree distance and reciprocal degree distance of open and closed fences,  $P_n \boxtimes K_2$  and  $C_n \boxtimes K_2$ , see Fig.4.




Fig. 4. Closed and open Fence graphs.

One can easily check that  $W(P_n) = \frac{n(n^2-1)}{6}$  and  $W(C_n) = \begin{cases} \frac{n^3}{8} & n \text{ is even} \\ \frac{n(n^2-1)}{8} & n \text{ is odd.} \end{cases}$

Similarly, we have  $DD(P_n) = \frac{1}{3}n(n-1)(2n-1)$  and  $DD(C_n) = 4W(C_n)$ .

One can observe that  $M_1(C_n) = 4n$ ,  $n \geq 3$ ,  $M_1(P_1) = 0$ , and  $M_1(P_n) = 4n-6$ ,  $n > 1$ . By direct calculations we obtain the Harary indices of  $P_n$  and  $C_n$

as follows.  $H(P_n) = n \left( \sum_{i=1}^n \frac{1}{i} \right) - n$  and  $H(C_n) = \begin{cases} n \left( \sum_{i=1}^{\frac{n}{2}} \frac{1}{i} \right) - 1, & \text{if } n \text{ is even} \\ n \left( \sum_{i=1}^{\frac{n-1}{2}} \frac{1}{i} \right), & \text{if } n \text{ is odd.} \end{cases}$

The following are the reciprocal degree distance of path and cycle on  $n$  vertices.  $RDD(P_n) = H(P_n) + 4 \left( \sum_{i=1}^{n-1} \frac{1}{i} \right) - \frac{3}{n-1}$  and  $RDD(C_n) = 4H(C_n)$ .

By using Corollaries 2.8 and 2.11, we obtain the exact formulae for degree distance and reciprocal degree distance of the following graphs.

EXAMPLE 2.12. (i)  $DD(P_n \boxtimes K_2) = \frac{4}{3} (5n^3 - 6n^2 + 31n - 24)$ .

(ii)  $DD(C_n \boxtimes K_2) = \begin{cases} 5n(n^2 + 2) & n \text{ is even} \\ 5n(n^2 + 1) & n \text{ is odd.} \end{cases}$

(iii)  $RDD(P_n \boxtimes K_2) = 16 \left( \sum_{i=1}^n \frac{1}{i} \right) + 32 \left( \sum_{i=1}^{n-1} \frac{1}{i} \right) - 6n - \frac{24}{n-1} - 8$ .

(iv)  $RDD(C_n \boxtimes K_2) = \begin{cases} 10n \left( 1 + 4 \sum_{i=1}^{\frac{n}{2}} \frac{1}{i} \right) - 40 & n \text{ is even} \\ 10n \left( 1 + 4 \sum_{i=1}^{\frac{n-1}{2}} \frac{1}{i} \right) & n \text{ is odd.} \end{cases}$

## ACKNOWLEDGMENTS

The authors wish to thank the referees for their recommendations to publish this article.

## REFERENCES

1. Y. Alizadeh, A. Iranmanesh, T. Doslic, Additively Weighted Harary Index of Some Composite Graphs, *Discrete Math.*, **313**, (2013), 26-34.
2. S. Chen, W. Liu, Extremal Modified Schultz Index of Bicyclic Graphs, *MATCH Commun. Math. Comput. Chem.*, **64**, (2010), 767-782.
3. J. Devillers, A. T. Balaban, *Topological Indices and Related Descriptors in QSAR and QSPR*, Gordon and Breach, Amsterdam, The Netherlands, 1999.
4. A. A. Dobrynin, A. A. Kochetova, Degree Distance of a Graph: a Degree Analogue of the Wiener Index, *J. Chem. Inf. Comput. Sci.*, **34**, (1994), 1082-1086.
5. T. Doslic, A. Graovac, F. Cataldo, O. Ori, Notes on Some Distance-Based Invariants for 2-Dimensional Square and Comb Lattices, *Iranian Journal of Mathematical Sciences and Informatics*, **2**(5), (2010), 61-68.
6. B. Furtula, I. Gutman, Z. Tomovic, A. Vesel, I. Pesek, Wiener-Type Topological Indices of Phenylenes, *Indian J. Chem.*, **41A**, (2002), 1767-1772.
7. I. Gutman, Selected Properties of the Schultz Molecular Topological Index, *J. Chem. Inf. Comput. Sci.*, **34**, (1994), 1087-1089.
8. I. Gutman, *O.E. Polansky, Mathematical Concepts in Organic Chemistry*, Springer-Verlag, Berlin, 1986.
9. I. Gutman, A Property of the Wiener Number and Its Modifications, *Indian J. Chem.*, **36A**, (1997), 128-132.
10. I. Gutman, A. A. Dobrynin, S. Klavzar, L. Pavlovic, Wiener-Type Invariants of Trees and Their Relation, *Bull. Inst. Combin. Appl.*, **40**, (2004), 23-30.
11. I. Gutman, D. Vidovic, L. Popovic, Graph Representation of Organic Molecules. Cayley's Plerograms vs. His Kenograms, *J. Chem. Soc. Faraday Trans.*, **94**, (1998), 857-860.
12. A. Hamzeh, A. Iranmanesh, S. Hossein-Zadeh, M.V. Diudea, Generalized Degree Distance of Trees, Unicyclic and Bicyclic Graphs, *Studia Ubb Chemia.*, LVII, **4**, (2012), 73-85.
13. A. Hamzeh, A. Iranmanesh, S. Hossein-Zadeh, Some Results on Generalized Degree Distance, *Open J. Discrete Math.*, **3**, (2013), 143-150.
14. B. Horoldagva, I. Gutman, On Some Vertex-Degree-based Graph Invariants, *MATCH Commun. Math. Comput. Chem.*, **65**, (2011), 723-730.
15. H. Hua, S. Zhang, On the Reciprocal Degree Distance of Graphs, *Discrete Appl. Math.*, **160**, (2012), 1152-1163.
16. O. Ivanciuc, T. S. Balaban, Reciprocal Distance Matrix Related Local Vertex Invariants and Topological Indices, *J. Math. Chem.*, **12**, (1993), 309-318.
17. A. Iranmanesh, Y. Aliazdeh, Computing Wiener Index of  $HAC5C7[p, q]$  Nanotubes by Gap program, *Iranian Journal of Mathematical Sciences and Informatics*, **1**(3), (2008), 1-12.
18. O. Khormali, A. Iranmanesh, I. Gutman, A. Ahamed, Generalized Schultz Index and Its Edge Versions, *MATCH Commun. Math. Comput. Chem.*, **64**, (2010), 783-798.
19. S. C. Li, X. Meng, Four Edge-Grafting Theorems on the Reciprocal Degree Distance of Graphs and Their Applications, *J. Comb. Optim.*, (2013), DOI 10.1007/s 10878-013-9649-1.
20. A. Mamut, E. Vumar, Vertex Vulnerability Parameters of Kronecker Products of Complete Graphs, *Inform. Process. Lett.*, **106**, (2008), 258-262.

21. K. Pattabiraman, M. Vijayaragavan, Reciprocal Degree Distance of Some Graph Operations, *Transactions on Combinatorics*, **2**, (2013), 13-24.
22. K. Pattabiraman, M. Vijayaragavan, Reciprocal Degree Distance of Product Graphs, *Discrete Appl. Math.*, **179**, (2014), 201-213.
23. G. F. Su, L.M. Xiong, X.F. Su, X.L. Chen, Some Results on the Reciprocal Sum-Degree Distance of Graphs, *J. Comb. Optim.*, (2013) DOI 10.1007/s 10878-013-9645-5.
24. I. Tomescu, Some Extremal Properties of the Degree Distance of a Graph, *Discrete Appl Math*, **98**, (1999), 159-163.
25. I. Tomescu, Properties of Connected Graphs Having Minimum Degree Distance, *Discrete Math*, **309**, (2008), 2745-2748.
26. H. Wang, L. Kang, Further Properties on the Degree Distance of Graphs, *J. Comb Optim.*, DOI 10.1007/s10878-014-9757-6.
27. H. Y. Zhu, D.J. Klenin, I. Lukovits, Extensions of the Wiener Number, *J. Chem. Inf. Comput. Sci.*, **36**, (1996), 420-428.