

Egoroff Theorem for Operator-Valued Measures in Locally Convex Cones

Davood Ayaseh^{a,*}, Asghar Ranjbari^b

^aDepartment of Pure Mathematics, Faculty of Mathematical Sciences,
University of Tabriz, Tabriz, Iran.

^bDepartment of Pure Mathematics, Faculty of Mathematical Sciences,
University of Tabriz, Tabriz, Iran.

E-mail: d_ayaseh@tabrizu.ac.ir

E-mail: ranjbari@tabrizu.ac.ir

ABSTRACT. In this paper, we define the almost uniform convergence and the almost everywhere convergence for cone-valued functions with respect to an operator valued measure. We prove the Egoroff theorem for \mathcal{P} -valued functions and operator valued measure $\theta : \mathfrak{R} \rightarrow \mathcal{L}(\mathcal{P}, \mathcal{Q})$, where \mathfrak{R} is a σ -ring of subsets of $X \neq \emptyset$, $(\mathcal{P}, \mathcal{V})$ is a quasi-full locally convex cone and $(\mathcal{Q}, \mathcal{W})$ is a locally convex complete lattice cone.

Keywords: Locally convex cones, Egoroff Theorem, Operator valued measure.

2000 Mathematics subject classification: 28A12, 46A03.

1. INTRODUCTION

The theory of locally convex cones as developed in [7] and [9] uses an order theoretical concept or convex quasi-uniform structure to introduce a topological structure on a cone. For recent researches see [1, 2, 3, 4, 8].

A *cone* is a set \mathcal{P} endowed with an addition and a scalar multiplication for nonnegative real numbers. The addition is assumed to be associative and commutative, and there is a neutral element $0 \in \mathcal{P}$. For the scalar multiplication the usual associative and distributive properties hold, that is $\alpha(\beta a) = (\alpha\beta)a$,

*Corresponding Author

$(\alpha + \beta)a = \alpha a + \beta a$, $\alpha(a + b) = \alpha a + \alpha b$, $1a = a$ and $0a = 0$ for all $a, b \in \mathcal{P}$ and $\alpha, \beta \geq 0$.

An *ordered cone* \mathcal{P} carries a reflexive transitive relation \leq such that $a \leq b$ implies $a + c \leq b + c$ and $\alpha a \leq \alpha b$ for all $a, b, c \in \mathcal{P}$ and $\alpha \geq 0$. The extended real numbers $\overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$ is a natural example of an ordered cone with the usual order and algebraic operations in $\overline{\mathbb{R}}$, in particular $0 \cdot (+\infty) = 0$.

A subset \mathcal{V} of the ordered cone \mathcal{P} is called an *abstract neighborhood system*, if the following properties hold:

- (1) $0 < v$ for all $v \in \mathcal{V}$;
- (2) for all $u, v \in \mathcal{V}$ there is a $w \in \mathcal{V}$ with $w \leq u$ and $w \leq v$;
- (3) $u + v \in \mathcal{V}$ and $\alpha v \in \mathcal{V}$ whenever $u, v \in \mathcal{V}$ and $\alpha > 0$.

For every $a \in \mathcal{P}$ and $v \in \mathcal{V}$ we define

$$v(a) = \{b \in \mathcal{P} | b \leq a + v\} \quad \text{resp.} \quad (a)v = \{b \in \mathcal{P} | a \leq b + v\},$$

to be a neighborhood of a in the upper, resp. lower topologies on \mathcal{P} . Their common refinement is called the symmetric topology generated by the neighborhoods $v^s(a) = v(a) \cap (a)v$. If we suppose that all elements of \mathcal{P} are bounded below, that is for every $a \in \mathcal{P}$ and $v \in \mathcal{V}$ we have $0 \leq a + \lambda v$ for some $\lambda > 0$, then the pair $(\mathcal{P}, \mathcal{V})$ is called a *full locally convex cone*. A *locally convex cone* $(\mathcal{P}, \mathcal{V})$ is a subcone of a full locally convex cone, not necessarily containing the abstract neighborhood system \mathcal{V} . For example, the extended real number system $\overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$ endowed with the usual order and algebraic operations and the neighborhood system $\mathcal{V} = \{\varepsilon \in \mathbb{R} | \varepsilon > 0\}$ is a full locally convex cone.

A subset B of the locally convex cone $(\mathcal{P}, \mathcal{V})$ is called *bounded below* whenever for every $v \in \mathcal{V}$ there is $\lambda > 0$, such that $0 \leq b + \lambda v$ for all $b \in B$.

For cones \mathcal{P} and \mathcal{Q} a mapping $T : \mathcal{P} \rightarrow \mathcal{Q}$ is called a *linear operator* if $T(a + b) = T(a) + T(b)$ and $T(\alpha a) = \alpha T(a)$ hold for all $a, b \in \mathcal{P}$ and $\alpha \geq 0$. If both \mathcal{P} and \mathcal{Q} are ordered, then T is called *monotone*, if $a \leq b$ implies $T(a) \leq T(b)$. If both $(\mathcal{P}, \mathcal{V})$ and $(\mathcal{Q}, \mathcal{W})$ are locally convex cones, the operator T is called (*uniformly*) *continuous* if for every $w \in \mathcal{W}$ one can find $v \in \mathcal{V}$ such that $T(a) \leq T(b) + w$ whenever $a \leq b + v$ for $a, b \in \mathcal{P}$.

A *linear functional* on \mathcal{P} is a linear operator $\mu : \mathcal{P} \rightarrow \overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$. The *dual cone* \mathcal{P}^* of a locally convex cone $(\mathcal{P}, \mathcal{V})$ consists of all continuous linear functionals on \mathcal{P} and is the union of all polars v° of neighborhoods $v \in \mathcal{V}$, where $\mu \in v^\circ$ means that $\mu(a) \leq \mu(b) + 1$, whenever $a \leq b + v$ for $a, b \in \mathcal{P}$. In addition to the given order \leq on the locally convex cone $(\mathcal{P}, \mathcal{V})$, the *weak preorder* \preceq is defined for $a, b \in \mathcal{P}$ by

$$a \preceq b \quad \text{if} \quad a \leq \gamma b + \varepsilon v$$

for all $v \in \mathcal{V}$ and $\varepsilon > 0$ with some $1 \leq \gamma \leq 1 + \varepsilon$ (for details, see [9], I.3). It is obviously coarser than the given order, that is $a \leq b$ implies $a \preceq b$ for $a, b \in \mathcal{P}$.

Given a neighborhood $v \in \mathcal{V}$ and $\varepsilon > 0$, the corresponding upper and lower relative neighborhoods $v_\varepsilon(a)$ and $(a)v_\varepsilon$ for an element $a \in \mathcal{P}$ are defined by

$$v_\varepsilon(a) = \{b \in \mathcal{P} \mid b \leq \gamma a + \varepsilon v \text{ for some } 1 \leq \gamma \leq 1 + \varepsilon\},$$

$$(a)v_\varepsilon = \{b \in \mathcal{P} \mid a \leq \gamma b + \varepsilon v \text{ for some } 1 \leq \gamma \leq 1 + \varepsilon\}.$$

Their intersection $v_\varepsilon^s(a) = v_\varepsilon(a) \cap (a)v_\varepsilon$ is the corresponding symmetric relative neighborhood. Suppose $v \in \mathcal{V}$. If we consider the abstract neighborhood system $\mathcal{V}_v = \{\alpha v : \alpha > 0\}$ on \mathcal{P} , then the corresponding upper (lower or symmetric) relative topology on \mathcal{P} is called *upper (lower or symmetric) relative v -topology*.

We shall say that a locally convex cone $(\mathcal{P}, \mathcal{V})$ is a *locally convex \vee -semilattice cone* if its order is antisymmetric and if for any two elements $a, b \in \mathcal{P}$ their supremum $a \vee b$ exists in \mathcal{P} and if

$$(\vee 1) (a + c) \vee (b + c) = a \vee b + c \text{ holds for all } a, b, c \in \mathcal{P},$$

$$(\vee 2) a \leq c + v \text{ and } b \leq c + w \text{ for } a, b, c \in \mathcal{P} \text{ and } v, w \in \mathcal{V} \text{ imply that } a \vee b \leq c + (v + w).$$

Likewise, $(\mathcal{P}, \mathcal{V})$ is a *locally convex \wedge -semilattice cone* if its order is antisymmetric and if for any two elements $a, b \in \mathcal{P}$ their infimum $a \wedge b$ exists in \mathcal{P} and if

$$(\wedge 1) (a + c) \wedge (b + c) = a \wedge b + c \text{ holds for all } a, b, c \in \mathcal{P},$$

$$(\wedge 2) c \leq a + v \text{ and } c \leq b + w \text{ for } a, b, c \in \mathcal{P} \text{ and } v, w \in \mathcal{V} \text{ imply that } c \leq a \wedge b + (v + w).$$

If both sets of the above conditions hold, then $(\mathcal{P}, \mathcal{V})$ is called a *locally convex lattice cone* (cf. [9]).

We shall say that a locally convex cone $(\mathcal{P}, \mathcal{V})$ is a *locally convex \vee^c -semilattice cone* if \mathcal{P} carries the weak preorder (that is the given order coincides with the weak preorder for the elements and the neighborhoods in \mathcal{P}), this order is antisymmetric and if

$$(\vee_1^c) \text{ every non-empty subset } A \subseteq \mathcal{P} \text{ has a supremum } \sup A \in \mathcal{P} \text{ and } \sup(A + b) = \sup A + b \text{ holds for all } b \in \mathcal{P},$$

$$(\vee_2^c) \text{ let } \emptyset \neq A \subseteq \mathcal{P}, b \in \mathcal{P} \text{ and } v \in \mathcal{V}. \text{ If } a \leq b + v \text{ for all } a \in A, \text{ then } \sup A \leq b + v.$$

Likewise, $(\mathcal{P}, \mathcal{V})$ is said to be a *locally convex \wedge^c -semilattice cone* if \mathcal{P} carries the weak preorder, this order is antisymmetric and if

$$(\wedge_1^c) \text{ every bounded below subset } A \subset \mathcal{P} \text{ has an infimum } \inf A \in \mathcal{P} \text{ and } \inf(A + b) = \inf A + b \text{ holds for all } b \in \mathcal{P},$$

$$(\wedge_2^c) \text{ let } A \subset \mathcal{P} \text{ be bounded below, } b \in \mathcal{P} \text{ and } v \in \mathcal{V}. \text{ If } b \leq a + v \text{ for all } a \in A, \text{ then } b \leq \inf A + v.$$

Combining both of the above notions, we shall say that a locally convex cone $(\mathcal{P}, \mathcal{V})$ is a *locally convex complete lattice cone* if \mathcal{P} is both a \vee^c -semilattice cone and a \wedge^c -semilattice cone.

As a simple, example the locally convex cone $(\overline{\mathbb{R}}, \mathcal{V})$, where $\overline{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$ and $\mathcal{V} = \{\varepsilon \in \mathbb{R} : \varepsilon > 0\}$, is a locally convex lattice cone and a locally convex complete lattice cone.

Suppose $(\mathcal{P}, \mathcal{V})$ is a locally convex complete lattice cone. A net $(a_i)_{i \in \mathcal{I}}$ in \mathcal{P} is called *bounded below* if there is $i_0 \in \mathcal{I}$ such that the set $\{a_i \mid i \geq i_0\}$ is bounded below. We define the superior and the inferior limits of a bounded below net $(a_i)_{i \in \mathcal{I}}$ in \mathcal{P} by

$$\liminf_{i \in \mathcal{I}} a_i = \sup_{i \in \mathcal{I}} (\inf_{k \geq i} a_k) \text{ and } \limsup_{i \in \mathcal{I}} a_i = \inf_{i \in \mathcal{I}} (\sup_{k \geq i} a_k).$$

If $\liminf_{i \in \mathcal{I}} a_i$ and $\limsup_{i \in \mathcal{I}} a_i$ coincide, then we denote their common value by $\lim_{i \in \mathcal{I}} a_i$ and say that the net $(a_i)_{i \in \mathcal{I}}$ is order convergent. A series $\sum_{i=1}^{\infty} a_i$ in $(\mathcal{P}, \mathcal{V})$ is said to be *order convergent* to $s \in \mathcal{P}$ if the sequence $s_n = \sum_{i=1}^n a_i$ is order convergent to s .

2. EGOROFF THEOREM FOR OPERATOR-VALUED MEASURES IN LOCALLY CONVEX CONES

The classical Egoroff theorem states that almost everywhere convergent sequences of measurable functions on a finite measure space converge almost uniformly. In this paper, we prove the Egoroff theorem for operator-valued measures in locally convex cones.

We shall say that a locally convex cone $(\mathcal{P}, \mathcal{V})$ is quasi-full if

(QF1) $a \leq b + v$ for $a, b \in \mathcal{P}$ and $v \in \mathcal{V}$ if and only if $a \leq b + s$ for some $s \in \mathcal{P}$ such that $s \leq v$, and

(QF2) $a \leq u + v$ for $a \in \mathcal{P}$ and $u, v \in \mathcal{V}$ if and only if $a \leq s + t$ for some $s, t \in \mathcal{P}$ such that $s \leq u$ and $t \leq v$.

The collection \mathfrak{A} of subsets of a set X is called a (weak) σ -ring whenever:

(R1) $\emptyset \in \mathfrak{A}$,

(R2) If $E_1, E_2 \in \mathfrak{A}$, then $E_1 \cup E_2 \in \mathfrak{A}$ and $E_1 \setminus E_2 \in \mathfrak{A}$,

(R3) If $E_n \in \mathfrak{A}$ for $n \in \mathbb{N}$ and $E_n \subseteq E$ for some $E \in \mathfrak{A}$, then $\bigcup_{n \in \mathbb{N}} E_n \in \mathfrak{A}$ (see [9]).

Any σ -algebra is a σ -ring and a σ -ring \mathfrak{A} is a σ -algebra if and only if $X \in \mathfrak{A}$.

However, we can associate with \mathfrak{A} in a canonical way the σ -algebra

$$\mathfrak{M}_{\mathfrak{A}} = \{A \subset X : A \cap E \in \mathfrak{A} \text{ for all } E \in \mathfrak{A}\}.$$

A subset A of X is said to be measurable whenever $A \in \mathfrak{M}_{\mathfrak{A}}$.

We consider the symmetric relative topology on \mathcal{P} . The function $f : X \rightarrow \mathcal{P}$ is measurable with respect to the σ -ring \mathfrak{A} if for every $v \in \mathcal{V}$,

(M₁) $f^{-1}(O) \cap E \in \mathfrak{A}$ for every open subset O of \mathcal{P} and every $E \in \mathfrak{A}$,

(M₂) $f(E)$ is separable in \mathcal{P} for every $E \in \mathfrak{A}$.

The operator-valued measures in locally convex cones have been defined in [9]. Let $(\mathcal{P}, \mathcal{V})$ be a quasi-full locally convex cone and let $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone. Let $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ denote the cone of all (uniformly)

continuous linear operators from \mathcal{P} to \mathcal{Q} . Recall from Section 3 in Chapter I from [9] that a continuous linear operator between locally convex cones is monotone with respect to the respective weak preorders. Because \mathcal{Q} carries its weak preorder, this implies monotonicity with respect to the given orders of \mathcal{P} and \mathcal{Q} as well. Let X be a set and \mathfrak{R} a σ -ring of subsets of X . An $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ -valued measure θ on \mathfrak{R} is a set function

$$E \rightarrow \theta_E : \mathfrak{R} \rightarrow \mathcal{L}(\mathcal{P}, \mathcal{Q})$$

such that $\theta_\emptyset = 0$ and

$$\theta_{(\bigcup_{n \in \mathbb{N}} E_n)} = \sum_{n \in \mathbb{N}} \theta_{E_n}$$

holds whenever the sets $E_n \in \mathfrak{R}$ are disjoint and $\bigcup_{n=1}^{\infty} E_n \in \mathfrak{R}$. Convergence for the series on the right-hand side is meant in the following way: For every $a \in \mathcal{P}$ the series $\sum_{n \in \mathbb{N}} \theta_{E_n}(a)$ is order convergent in \mathcal{Q} . We note that the order convergence is implied by convergence in the symmetric relative topology.

Let $(\mathcal{P}, \mathcal{V})$ be a quasi-full locally convex cone and let $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone. Suppose θ is a fixed $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ -valued measure on \mathfrak{R} . For a neighborhood $v \in \mathcal{V}$ and a set $E \in \mathfrak{R}$, semivariation of θ is defined as follows:

$$|\theta|(E, v) = \sup \left\{ \sum_{i \in \mathbb{N}} \theta_{E_i}(s_i) : s_i \in \mathcal{P}, s_i \leq v, E_i \in \mathfrak{R} \text{ disjoint subsets of } E \right\}.$$

It is proved in Lemma 3.3 chapter II from [9], that if $v \in \mathcal{P}$, then $|\theta|(E, v) = \theta_E(v)$.

Proposition 2.1. *Let $(\mathcal{P}, \mathcal{V})$ be a quasi-full locally convex cone, $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone and θ be a fixed $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ -valued measure on \mathfrak{R} .*

- (a) *If for $E \in \mathfrak{R}$, $\theta_E = 0$, then for every $v \in \mathcal{V}$, $|\theta|(E, v) = 0$,*
 (b) *If for every $v \in \mathcal{V}$, $|\theta|(E, v) = 0$, then $\theta_E(a) = 0$ for every bounded element a of \mathcal{P} .*

Proof. For (a), let $\theta_E = 0$ and $F_1, \dots, F_n, n \in \mathbb{N}$ be a partition of E . Then for $0 \leq s_i \leq v, i = 1, \dots, n$, we have $0 \leq \theta_{F_i}(s_i) \leq \theta_E(s_i) = 0$. Since the order of \mathcal{Q} is antisymmetric, for every $i \in \{1, \dots, n\}$, we have $\theta_{F_i}(s_i) = 0$. Then $|\theta|(E, v) = 0$.

For (b), let $a \in \mathcal{P}$ and for every $v \in \mathcal{V}$, $|\theta|(E, v) = 0$. Since a is bounded, for $v \in \mathcal{V}$, there is $\lambda > 0$ such that $0 \leq a + \lambda v$ and $a \leq \lambda v$. Now we have $0 \leq \theta_E(a) + |\theta|(E, \lambda v)$ and $\theta_E(a) \leq |\theta|(E, \lambda v)$ by Lemma II,3.4 of [9]. This shows that $0 \leq \theta_E(a)$ and $\theta_E(a) \leq 0$. Since the order of \mathcal{Q} is antisymmetric, we have $\theta_E(a) = 0$. \square

Corollary 2.2. *Let $(\mathcal{P}, \mathcal{V})$ be a quasi-full locally convex cone, $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone and θ be a fixed $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ -valued measure on*

\mathfrak{A} . If all elements of \mathcal{P} are bounded, then for $E \in \mathfrak{A}$, $\theta_E = 0$ if and only if $|\theta|(E, v) = 0$ for all $v \in \mathcal{V}$.

Definition 2.3. Let \mathfrak{A} be a σ -ring of subsets of X . The set $A \in \mathfrak{A}$ is said to be of positive v -semivariation of the measure θ if $|\theta|(A, v) > 0$. Also, we say that the set A has bounded v -semivariation of the measure θ , if $|\theta|(A, v)$ is bounded in $(\mathcal{Q}, \mathcal{W})$.

Definition 2.4. Let θ be an operator-valued measure on X . We shall say that θ is *generalized strongly v -continuous* (GS_v -continuous, for short) if for every set of bounded v -semivariation $E \in \mathfrak{A}$ and every monotone sequence of sets $(E_n)_{n \in \mathbb{N}} \in \mathfrak{A}$, $E_n \subset E$, $n \in \mathbb{N}$ the following holds

$$\lim_{n \in \mathbb{N}} |\theta|(E_n, v) = |\theta|(\lim_{n \in \mathbb{N}} E_n, v) \quad v \in \mathcal{V},$$

where the limit in the left hand side of the equality means convergence with respect to the symmetric relative topology of $(\mathcal{Q}, \mathcal{W})$.

EXAMPLE 2.5. Let $X = \mathbb{N} \cup \{+\infty\}$ and $\mathcal{P} = \mathcal{Q} = \bar{\mathbb{R}}$. We consider on $\bar{\mathbb{R}}$ the abstract neighborhood system $\mathcal{V} = \{\varepsilon \in \mathbb{R} : \varepsilon > 0\}$. Then $\mathcal{L}(\mathcal{P}, \mathcal{Q})$ contains all nonnegative reals and the linear functional $\bar{0}$ acting as

$$\bar{0}(x) = \begin{cases} +\infty & x = +\infty \\ 0 & \text{else.} \end{cases}$$

We set $\mathfrak{A} = \{E \subset X : E \text{ is finite}\}$. Then \mathfrak{A} is a σ -ring on X . We define the set function θ on \mathfrak{A} as following: for $x \in X$, $\theta_\emptyset = 0$, $\theta_{\{n\}}(x) = nx$ for $n \in \mathbb{N}$ and $\theta_{\{+\infty\}}(x) = \bar{0}(x)$. For $E = \{a_1, \dots, a_n\} \in \mathfrak{A}$, $n \in \mathbb{N}$, we define $\theta_E(x) = \sum_{i=1}^n \theta_{\{a_i\}}(x)$ for $x \in X$. Then θ is clearly an operator-valued measure on \mathfrak{A} . For $n \in \mathbb{N}$ and $\varepsilon > 0$, we have $|\theta|(\{n\}, \varepsilon) = \theta_{\{n\}}(\varepsilon) = n\varepsilon$ and $|\theta|(\{+\infty\}, \varepsilon) = \theta_{\{+\infty\}}(\varepsilon) = \bar{0}(\varepsilon) = 0$. Therefore each $E \in \mathfrak{A}$ has finite ε -semivariation for all $\varepsilon > 0$. Let $E \in \mathfrak{A}$. If $(E_n)_{n \in \mathbb{N}} \subset \mathfrak{A}$ is a monotone sequence of subsets of E such that $\lim_{n \in \mathbb{N}} E_n = F$, then there is $n_0 \in \mathbb{N}$ such that $E_n = F$ for all $n \geq n_0$. Then θ is clearly GS_ε -continuous for each $\varepsilon > 0$.

Definition 2.6. A sequence $(f_n)_{n \in \mathbb{N}}$ of measurable functions is said to be θ -almost uniformly convergent to a measurable function f on $E \in \mathfrak{A}$ if for every $\varepsilon > 0$, $w \in \mathcal{W}$ and $v \in \mathcal{V}$ there exists a subset $F = F(\varepsilon, v, w)$ of E and $n_0 \in \mathbb{N}$ such that for every $n > n_0$,

$$f_n(x) \in v_\varepsilon^s(f(x)) \text{ and } |\theta|(F, v) \in w_\varepsilon^s(0),$$

for all $x \in E \setminus F$.

Theorem 2.7 (Egoroff Theorem). Let \mathfrak{A} be a σ -ring of subsets of X , $(\mathcal{P}, \mathcal{V})$ be a full locally convex cone and $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone. For $v \in \mathcal{V}$, suppose $\theta : \mathfrak{A} \rightarrow \mathcal{L}(\mathcal{P}, \mathcal{Q})$ is a GS_v -continuous operator valued measure, and $E \in \mathfrak{A}$ has bounded v -semivariation. If $f : X \rightarrow \mathcal{P}$ is a measurable function, and $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ is a sequence of measurable

functions, such that for every $t \in E$, $f_n(t) \rightarrow f(t)$ with respect to the symmetric relative v -topology of $(\mathcal{P}, \mathcal{V})$, then $(f_n)_{n \in \mathbb{N}}$ is θ -almost uniformly convergent to f on E , with respect to the symmetric relative v -topology of $(\mathcal{P}, \mathcal{V})$.

Proof. We identify $v \in \mathcal{V}$ with the constant function $x \rightarrow v$ from X into \mathcal{P} . For $m, n \in \mathbb{N}$, we set

$$B_n^m = \bigcap_{i=n}^{\infty} \{x \in E : f_i(x) \preceq_v f(x) + \frac{1}{m}v \text{ and } f(x) \preceq_v f_i(x) + \frac{1}{m}v\}.$$

For every $n, m \in \mathbb{N}$ we have $B_n^m \in \mathfrak{R}$ by Theorem II.1.6 from [9]. Clearly, $B_n^m \subset B_{n+1}^m$ for all $n, m \in \mathbb{N}$. We claim that $E = \bigcup_{n=1}^{\infty} B_n^m$. Let $x \in E$ and $m \in \mathbb{N}$. Then $(f_n(x))_{n \in \mathbb{N}}$ is convergent to $f(x)$ with respect to the symmetric relative v -topology. This shows that for each $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$ such that $f_n(x) \in (\frac{1}{m}v)_{\varepsilon}^s(f(x))$ for all $n \geq n_0$. Therefore $f_n(x) \leq \gamma f(x) + \varepsilon(\frac{1}{m}v)$ and $f_n(x) \leq \gamma f(x) + \varepsilon(\frac{1}{m}v)$ for all $n \geq n_0$ and some $1 \leq \gamma \leq 1 + \varepsilon$. This yields that $f_n(x) \leq \gamma f(x) + (1 + \varepsilon)(\frac{1}{m}v)$ and $f_n(x) \leq \gamma f(x) + (1 + \varepsilon)(\frac{1}{m}v)$ for all $n \geq n_0$ and some $1 \leq \gamma \leq 1 + \varepsilon$. Now Lemma I.3.1 from [9] shows that $f_n(x) \preceq_v f(x) + \frac{1}{m}v$ and $f(x) \preceq_v f_n(x) + \frac{1}{m}v$ for all $n \geq n_0$. Thus $x \in B_{n_0}^m$.

Then $(E \setminus B_n^m)_{n \in \mathbb{N}}$ is a decreasing sequence of subsets of E , such that $\lim_{n \rightarrow \infty} E \setminus B_n^m = \emptyset$. Therefore for every $m \in \mathbb{N}$, $|\theta|(E \setminus B_n^m, v)$ is convergent to $|\theta|(\emptyset, v) = 0$ with respect to the symmetric relative topology of $(\mathcal{Q}, \mathcal{W})$ by the assumption. For $\varepsilon > 0$ and $m \in \mathbb{N}$ we choose n_m such that $|\theta|(E \setminus B_{n_m}^m, v) \leq \frac{\varepsilon}{2^m}w$. We set

$$F = \bigcup_{m=1}^{\infty} E \setminus B_{n_m}^m.$$

Then we have

$$|\theta|(F, v) \leq \sum_{m=1}^{\infty} |\theta|(B_{n_m}^m, v) \leq \sum_{m=1}^{\infty} \frac{\varepsilon}{2^m}w = \varepsilon w.$$

Also, we have $0 \leq |\theta|(F, v) + \varepsilon w$. Then $|\theta|(F, v) \in w_{\varepsilon}^s(0)$.

Now, we show that the convergence on $E \setminus F$ is uniform. Let $\delta > 0$. There is $k \in \mathbb{N}$ such that $\frac{2}{k} + \frac{1}{k^2} \leq \delta$. We have

$$E \setminus F = E \setminus \left(\bigcup_{m=1}^{\infty} E \setminus B_{n_m}^m \right) = \bigcap_{m=1}^{\infty} B_{n_m}^m \subset B_{n_k}^k$$

Now for each $n \geq n_k$ and every $x \in E \setminus F$ we have $f_n(x) \preceq_v f(x) + \frac{1}{k}v$ and $f(x) \preceq_v f_n(x) + \frac{1}{k}v$. The definition of \preceq_v shows that for $\varepsilon = \frac{1}{k}$ there is $1 \leq \gamma \leq$

$1 + \frac{1}{k}$ such that $f_n(x) \leq \gamma(f(x) + \frac{1}{k}v) + \frac{1}{k}v$ and $f(x) \leq \gamma(f_n(x) + \frac{1}{k}v) + \frac{1}{k}v$. Therefore $f_n(x) \leq \gamma f(x) + (\frac{2}{k} + \frac{1}{k^2})v \leq \gamma f(x) + \delta v$ and $f(x) \leq \gamma f_n(x) + (\frac{2}{k} + \frac{1}{k^2})v \leq \gamma f_n(x) + \delta v$. Since $1 \leq \gamma \leq 1 + \frac{1}{k} \leq 1 + \frac{2}{k} + \frac{1}{k^2} \leq 1 + \delta$, we realize that $(f_n)_{n \in \mathbb{N}}$ is uniformly convergent to f on $E \setminus F$, with respect to the symmetric relative topology. \square

Remark 2.8. If in the assumptions of Theorem 2.7, $(\mathcal{P}, \mathcal{V})$ is a quasi-full locally convex cone, then the theorem holds again. In fact every quasi-full locally convex cone can be embedded in a full locally convex cone as elaborated in ([9], I, 6.2).

Definition 2.9. We say that a sequence $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ of measurable functions is θ -almost everywhere convergent (with respect to the symmetric topology of $(\mathcal{P}, \mathcal{V})$) to f , if the set $\{x \in X : f_n(x) \not\rightarrow f(x)\}$ is contained in a subset E of X with $\theta_E = 0$.

Definition 2.10. Let $v \in \mathcal{V}$. We say that the sequence $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ of measurable functions is $|\theta|_v$ -almost everywhere convergent (with respect to symmetric topology of $(\mathcal{P}, \mathcal{V})$) to f , if the set $\{x \in X : f_n(x) \not\rightarrow f(x)\}$ is contained in a subset E of X with $|\theta|(E, v) = 0$.

Lemma 2.11. Let \mathfrak{A} be a σ -ring of subsets of X , $(\mathcal{P}, \mathcal{V})$ be a full locally convex cone and $(\mathcal{Q}, \mathcal{W})$ be a locally convex complete lattice cone. Then

- (a) θ -almost everywhere convergence implies $|\theta|_v$ -almost everywhere convergence for each $v \in \mathcal{V}$.
- (b) If all elements of $(\mathcal{P}, \mathcal{V})$ are bounded and a sequence $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ is $|\theta|_v$ -almost everywhere convergent to f for each $v \in \mathcal{V}$, then $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ is θ -almost everywhere convergent to f .

Proof. The assertions are proved by the help of Proposition 2.1. \square

Theorem 2.12. If in the Egoroff theorem (2.7), $f_n \rightarrow f$, θ -almost everywhere or $|\theta|_v$ -almost everywhere, then the assertion of theorem holds.

Proof. Suppose $f_n \rightarrow f$, θ -almost everywhere, then there is a subset A of E , which is contained in some $B \in \mathfrak{A}$ with $\theta_B = 0$. Now $E \setminus B \in \mathfrak{A}$ and it has bounded v -semivariation. We apply the theorem 2.7 for $E \setminus B$ and obtain a subset F satisfying in definition 2.6. Now clearly f_n is θ -almost uniformly convergent to f on $E \setminus (F \cap B)$. A similar argument yields our claim for $|\theta|_v$ -almost everywhere convergence. \square

Theorem 2.13. Let the symmetric relative w -topology of $(\mathcal{Q}, \mathcal{W})$ be Hausdorff for each $w \in \mathcal{W}$ and let $(f_n : X \rightarrow \mathcal{P})_{n \in \mathbb{N}}$ be a sequence of measurable functions which converges to f , θ -almost uniformly on $E \in \mathfrak{A}$. Then $\{f_n\}_{n \in \mathbb{N}}$ is $|\theta|_v$ -almost everywhere convergent to f for each $v \in \mathcal{V}$.

Proof. For each $n \in \mathbb{N}$, $v \in \mathcal{V}$ and $w \in \mathcal{W}$ there is $F_n = F_n(v, w) \in \mathfrak{A}$ such that $F_n \subset E$ and $|\theta|(F_n, v) \in w_{\frac{1}{n}}^s(0)$ and (f_n) is convergent to f on $E \setminus F_n$. Now, we set $F = \bigcap_{n=1}^{\infty} F_n$. Since $(\mathcal{Q}, \mathcal{W})$ is separated, we have $|\theta|(F, v) = 0$. Clearly, $(f_n(x))_{n \in \mathbb{N}}$ is convergent to $f(x)$ for each $x \in E \setminus F = \bigcup_{n=1}^{\infty} E \setminus F_n$. \square

ACKNOWLEDGMENTS

The authors would like to thank the referees for their useful comments.

REFERENCES

1. D. Ayaseh, A. Ranjbari, Bornological Convergence in Locally Convex Cones, *Mediterr. J. Math.*, **13**(4), (2016), 1921-1931.
2. D. Ayaseh, A. Ranjbari, Bornological Locally Convex Cones, *Le Matematiche*, **69**(2), (2014), 267-284.
3. D. Ayaseh, A. Ranjbari, Locally Convex Quotient Lattice Cones, *Math. Nachr.*, **287**(10), (2014), 1083-1092.
4. D. Ayaseh, A. Ranjbari, Some Notes on Bornological and Nonbornological Locally Convex Cones, *Le Matematiche*, **70**(2), (2015), 235-241.
5. N. Eftekhari, The Basic Theorem and its Consequences, *Iranian Journal of Mathematical Sciences and Informatics*, **4**(1), (2009), 27-35.
6. P. R. Halmos, *Measure Theory*, Van Nostrand, Princeton, N. J., 1950. MR 11, 504.
7. K. Keimel, W. Roth, Ordered Cones and Approximation, *Lecture Notes in Mathematics*, **1517**, 1992, Springer Verlag, Heidelberg-Berlin-New York.
8. A. Ranjbari, Strict Inductive Limits in Locally Convex Cones, *Positivity*, **15**(3), (2011), 465-471.
9. W. Roth, Operator-valued Measures and Integrals for Cone-valued Functions, *Lecture Notes in Mathematics*, **1964**, 2009, Springer Verlag, Heidelberg-Berlin-New York.
10. A. Taghavi, R. Hosseinzadeh, Uniform Boundedness Principle for Operators on Hypervector Spaces, *Iranian Journal of Mathematical Sciences and Informatics*, **7**(2), (2012), 9-16.