Iranian Journal of Mathematical Sciences and Informatics

Vol. 20, No. 2 (2025), pp 17-29 DOI: 10.61186/ijmsi.20.2.17

Hypervaluations on Hyperfields and Ordered Canonical Hypergroups

Alessandro Linzi a* , Hanna Stojałowska b

 ^aCentre for Information Technologies and Applied Mathematics University of Nova Gorica, Vipavska 13, SI-5000 Nova Gorica, Slovenia
 ^bDepartment of Mathematics West Pomeranian University of Technology, Piastów 48, PL 70-311 Szczecin, Poland

E-mail: linzi.alessandro@gmail.com
E-mail: hanna.stojalowska@phd.usz.edu.pl

ABSTRACT. We study the concept of hypervaluations on hyperfields. The main aim of this paper is to show that any hypervaluation from a hyperfield onto an ordered canonical hypergroup is the composition of a hypervaluation onto an ordered abelian group (which induces the same valuation hyperring) and an order preserving homomorphism of hypergroups. This implies that the generalization of hypervaluations proposed in [13] can be reduced to the original definition of hypervaluation given by Davvaz and Salasi in [4]. Moreover, we provide counterexamples to some assertions which appear in [13].

Keywords: Hypergroup, Hyperfield, Hyperring, Hypervaluation, Ordered canonical hypergroup.

2000 Mathematics subject classification: Primary: 20N20, 06F99, Secondary: 13A18.

Received 16 January 2021; Accepted 07 February 2022 © 2025 Academic Center for Education, Culture and Research TMU

^{*}Corresponding Author

1. Introduction

The theory of hyperrings and hyperfields has its origins in the paper of Krasner [7]. More recently, several authors studied their theory from various points of view. For instance, J. Jun, in [6], studies algebraic geometry over hyperrings and gives the definition of hyperideals in a hyperring. The reader may consult the reference list of this paper to gain an overview over recent developments in the theory of hyperfields and their important applications.

M. Marshall in [11] studies the definitions of orderings and of positive cones in hyperfields. It was already mentioned in that paper that the two concepts are not equivalent as in the classical theory of ordered fields. In fact, we show in Example 2.15 that the natural generalization of the construction which for ordered fields shows this equivalence, does not work in the case of hyperfields.

In [4] B. Davvaz and A. Salasi deal with hypervaluations on a hyperring onto an ordered abelian group, and also J. Lee in [10] works with valued hyperfields. These two papers present different approaches in defining a hypervaluation on a hyperfield, which both appear to be interesting and well chosen.

The interest of the authors for the concepts of hypervaluations and hyperfields arises mainly from the fact that there are hyperfields related to valued fields which inherit a hypervaluation in a natural way. In [9] amc-structures are introduced by F.-V. Kuhlmann in order to study criteria for valued fields to be isomorphic or to be elementarily equivalent. In [5] Flenner introduced RV-structures which are an improvement of amc-structures and are nothing but hyperfields presented in a different language, i.e., with a ternary relation replacing and encoding the multi-valued addition. Therefore, as can be expected, also amc and RV structures implicitly have (hyper)valuations. These can be used to study them and the corresponding hyperfields further.

In this paper we study a possible definition of hypervaluation, introduced in [13]. What is proposed there is a generalization of the definition of Davvaz and Salasi where the value set is allowed to be an ordered canonical hypergroup (see Definition 2.10). Here the domain is always assumed to be a hyperfield. We show in our main result that even though this definition is proper, i.e., nontrivial examples can be found, it is not of much interest since such a hypervaluation can always be decomposed into an order preserving homomorphism of hypergroups and a hypervaluation onto an ordered abelian group (in the sense of Davvaz and Salasi) which, moreover, induces the same valuation hyperring.

In the paper [13], a lot of constructions are proposed without details; it turns out that they are not always possible to carry out. In the present article we also provide counterexamples to justify this last assertion.

2. Ordered Canonical Hypergroups

Definition 2.1 ([1]). Let H be a nonempty set and $\mathcal{P}^*(H)$ the family of nonempty subsets of H. A hyperoperation * is a function which associates with every pair $(x, y) \in H \times H$ an element of $\mathcal{P}^*(H)$, denoted by x * y.

A hypergroupoid is a nonempty set H with a hyperoperation $*: H \times H \to \mathcal{P}^*(H)$. For $x \in H$, $A, B \subseteq H$ we set

$$A*B=\bigcup_{a\in A,b\in B}a*b,$$

$$A * x = A * \{x\}$$
and $x * A = \{x\} * A$.

In 1934 the concept of a hypergroup was defined by F. Marty in [12] to be a nonempty set H with an associative hyperoperation (see Definition 2.2 below) such that x * H = H * x = H for all $x \in H$. A special class of hypergroups, which will be of interest for us, is the following:

Definition 2.2 ([1]). A canonical hypergroup is a tuple (H, *, e), where (H, *) is a hypergroupoid and e is an element of H such that the following axioms hold:

- (H1) the hyperoperation * is associative, i.e., (x*y)*z = x*(y*z) for all $x, y, z \in H$,
- (H2) x * y = y * x for all $x, y \in H$,
- (H3) for every $x \in H$ there exists a unique $x' \in H$ such that $e \in x * x'$ (the element x' will be denoted by x^{-1}),
- (H4) $z \in x * y$ implies $y \in x^{-1} * z$ for all $x, y, z \in H$.

Remark 2.3. A canonical hypergroup is a hypergroup in the sense of Marty. Fix $a \in H$ and take $x \in H * a$. Then there exist $h \in H$ such that $x \in h * a \subseteq H$, showing that $H * a \subseteq H$. For the other inclusion, take $x \in H$, then

$$x \in x * e \subseteq x * (a^{-1} * a) = (x * a^{-1}) * a,$$

so there exist $h \in x * a^{-1} \subseteq H$ such that $x \in h * a \subseteq H * a$.

Remark 2.4. In [1] the definition of a canonical hypergroup also requires explicitly that $x*e = \{x\}$ for all $x \in H$. However, we note that this axiom follows from (H3) and (H4). Indeed, suppose that $y \in x*e$ for some $x, y \in H$. Then $e \in x^{-1} * y$ by (H4). Now y = x follows from the uniqueness required in (H3).

Remark 2.5. Note that an abelian group G is not a priori a hypergroup, because the operation on G is not a hyperoperation, as it takes values in G and not in $\mathcal{P}^*(G)$. But it can be turned into a hypergroup by setting $a*b := \{ab\}$. In other words, we can turn an abelian group into a hypergroup by identifying each element a of G with the singleton $\{a\}$.

EXAMPLE 2.6. Consider the set $H := \{-1, 0, 1\}$ with a hyperoperation * defined as follows:

$$(-1) * (-1) = (-1) * 0 = 0 * (-1) = \{-1\}$$
$$0 * 0 = \{0\}$$
$$1 * 1 = 1 * 0 = 0 * 1 = \{1\}$$
$$1 * (-1) = (-1) * 1 = \{-1, 0, 1\}.$$

Then (H, *, 0) is a canonical hypergroup, called the *sign hypergroup*. As the reader may check, we have that $1^{-1} = -1$, $(-1)^{-1} = 1$, $0^{-1} = 0$.

The next example can be found in [8].

EXAMPLE 2.7. Let R be a ring and G a normal subgroup of its multiplicative semigroup. Consider the following equivalence relation \sim on R: $a \sim b$ if and only if there exist $g, h \in G$ s.t. ag = bh. The equivalence class of $a \in R$ is

$$aG := \{ ag \mid g \in G \}.$$

It is possible to define a hyperoperation on R/G in the following way:

$$aG + bG := \{(ag + bh)G \mid g, h \in G\}.$$

Then $(R/G, +, \{0_R\})$ is a canonical hypergroup. Indeed, the associative law follows from the same law in R, as well as commutativity. The unique inverse of aG is (-a)G. Indeed,

$$aG + (-a)G = \{(ag - ah)G \mid g, h \in G\} \ni (a - a)G = 0_R G,$$

moreover, if

$$0_R G \in aG + bG = \{(ag + bh)G \mid g, h \in G\},\$$

then there exist $g, h \in G$ such that $0_R = ag + bh$. Multiplying by g^{-1} , we obtain that $-a = bhg^{-1}$ and so (-a)G = bG.

Assume now that $cG \in aG + bG$. We wish to show that $bG \in (-a)G + cG$. We have

$$cG \in \{(ag + bh)G \mid g, h \in G\},\$$

so there exist $g, h \in G$ such that c = ag + bh. Multiplying by h^{-1} , we obtain $b = -agh^{-1} + ch^{-1}$, so $bG \in \{((-a)g' + ch')G \mid g', h' \in G\} = (-a)G + cG$.

Remark 2.8. We note that the sign hypergroup can be obtained as a quotient in the way described in the previous example. Take $R = \mathbb{R}$ and $G = \dot{\mathbb{R}}^2$, where $\dot{\mathbb{R}}^2$ denotes the set of non-zero squares in \mathbb{R} . The result follows from the fact that every non-zero real number is either a square or the opposite of a square.

In [13] the set $\mathbb{Z}/\mathbb{N} = \{a\mathbb{N} \mid a \in \mathbb{Z}\}$ is considered, with the hyperoperation defined as follows:

$$a\mathbb{N} * b\mathbb{N} = \{c\mathbb{N} \mid c \in a\mathbb{N} + b\mathbb{N}\}.$$

This construction does not give a canonical hypergroup in the sense of Definition 2.2, as is shown in the next example. The reason is that \mathbb{N} is not a multiplicative subgroup of the multiplicative semigroup of \mathbb{Z} .

EXAMPLE 2.9. Consider the tuple $(\mathbb{Z}/\mathbb{N}, *, 0\mathbb{N})$. We observe that $0\mathbb{N} \in 1\mathbb{N}*-k\mathbb{N}$ for every $k \in \mathbb{N}$. Hence axiom (H3) is not fulfilled.

Definition 2.10. An ordered canonical hypergroup is a tuple $(H, *, e, \leq)$, where (H, *, e) is a canonical hypergroup and \leq is a partial order such that

$$a < b \Longrightarrow a * c \nearrow b * c$$
 (2.1)

for all $a, b, c \in H$; here, if $A, B \subseteq H$, then $A \nearrow B$ means that for all $b \in B$ there exists $a \in A$ such that $a \leq b$. If for each $a, b \in H$ either $a \leq b$ or $b \leq a$, then we call \leq an ordering or a linear order relation on H.

Remark 2.11. One should not use " \leq " on the subsets as the relation will not be antisymmetric. For example if H contains a smallest element b, then any two subsets A, B, which both contain b will satisfy $A \nearrow B$ and $B \nearrow A$ without necessarily being equal.

Remark 2.12. It should be noted that some authors (for instance see [13]) in defining $A \nearrow B$ require that for all $a \in A$ there exists $b \in B$ and for all $b \in B$ there exists $a \in A$ such that $a \leq b$. Others (for instance see [3]) instead require only that for all $a \in A$ there exists $b \in B$ such that $a \leq b$. The relations between all these definitions still have to be investigated. However, in what follows we will just use the concept introduced in Definition 2.10.

Lemma 2.13. Let $(H, *, e, \leq)$ be an ordered canonical hypergroup. Take $a, b, x, y \in H$ and $B \subseteq H$.

- 1) If $\{a\} \nearrow B$, then a < b for all $b \in B$.
- 2) If x > e, then $x^{-1} < e$.
- 3) If $x \ge e$ and $y \ge e$, then $b \ge e$ for all $b \in x * y$.
- 4) If x > e and $y \ge e$, then b > e for all $b \in x * y$.

Proof. 1): This follows from the definition of " \nearrow " and the fact that $\{a\}$ contains only a.

- 2): By condition (2.1), $\{x^{-1}\}=e*x^{-1}\nearrow x*x^{-1}$, so $x^{-1}\le b$ for every $b\in x*x^{-1}$ by part 1). As $e\in x*x^{-1}$, we obtain that $x^{-1}\le e$. Now $x^{-1}=e$ is impossible because otherwise, $e\in x*x^{-1}=\{x\}$, which implies that x=e in contradiction to our assumption that x>e.
- 3): If $x \ge e$ and $y \ge e$, then $\{y\} = e * y \nearrow x * y$, so $e \le y \le b$ for every $b \in x * y$ by part 1).
- 4): If x > e and $y \ge e$, then $b \ge e$ for all $b \in x * y$ by part 3). However, $e \notin x * y$ since otherwise, $y = x^{-1}$ which by part 2) yields that y < e, in contradiction to our assumption that $y \ge e$. Hence $b \ne e$ and consequently, b > e.

In [11] the definition of a positive cone is given for hyperfields. Let us now define this concept for canonical hypergroups.

Definition 2.14 ([13]). A subset P of a canonical hypergroup (H, *, e) is called a *positive cone* if the following axioms hold:

- (P1) $P \cap -P = \{e\},\$
- (P2) $P * P \subseteq P$,
- (P3) $P \cup -P = H$.

In the theory of ordered abelian groups, the existence of a positive cone is equivalent to the existence of an ordering and there is a one to one correspondence between them. It was already mentioned in [11] that this, in general, is no more true in the case of hyperfields, and the argument holds for hypergroups as well. In [13] it is claimed that one can always construct an ordering from a positive cone by setting:

$$x \le y \iff (y * x^{-1}) \cap P \ne \emptyset. \tag{2.2}$$

In the following example we show that this construction is not always possible.

EXAMPLE 2.15. Take $H := \mathbb{Q}/\dot{\mathbb{Q}}^2$, where $\dot{\mathbb{Q}}^2$ denotes the set of nonzero squares in \mathbb{Q} . We see that H is a canonical hypergroup with the hyperoperation defined as:

$$a\dot{\mathbb{Q}}^2 * b\dot{\mathbb{Q}}^2 = \{c\dot{\mathbb{Q}}^2 \mid c \in a\dot{\mathbb{Q}}^2 + b\dot{\mathbb{Q}}^2\}$$

(see also Example 2.7). Observe that the set $P = \{a\dot{\mathbb{Q}}^2 \mid a \in \mathbb{Q}^+\} \cup \{0\}$ fulfils the conditions of the definition of a positive cone. However, the relation defined in (2.2) is not an ordering on H. Indeed, observe that $5\dot{\mathbb{Q}}^2 \in (2\dot{\mathbb{Q}}^2*(3\dot{\mathbb{Q}}^2)^{-1}) \cap P$, so, in particular, $(2\dot{\mathbb{Q}}^{*2}*(3\dot{\mathbb{Q}}^2)^{-1}) \cap P \neq \emptyset$, which means that $3\dot{\mathbb{Q}}^2 \leq 2\dot{\mathbb{Q}}^2$. On the other hand, $1\dot{\mathbb{Q}}^2 \in (3\dot{\mathbb{Q}}^2*(2\dot{\mathbb{Q}}^2)^{-1}) \cap P$, so $(3\dot{\mathbb{Q}}^2*(2\dot{\mathbb{Q}}^2)^{-1}) \cap P \neq \emptyset$, which means that $2\dot{\mathbb{Q}}^2 \leq 3\dot{\mathbb{Q}}^2$. Clearly, $2\dot{\mathbb{Q}}^2 \neq 3\dot{\mathbb{Q}}^2$, so the relation \leq is not antisymmetric.

However, there exist hypergroups in which positive cone and orderings behave as in the classical case. The simplest example is the sign hypergroup:

EXAMPLE 2.16. Consider the sign hypergroup $H = \{-1, 0, 1\}$ with positive cone $P = \{0, 1\}$. We define an order relation \leq on H as follows: $-1 \leq 0 \leq 1$. We leave it to the reader to show that \leq is a linear order on H as in Definition 2.10. This ordering corresponds to P in the way described in (2.2). Indeed, $(1*0^{-1}) \cap P = \{1\}, (0*(-1)^{-1}) \cap P = \{1\}, (1*(-1)^{-1}) \cap P = \{1\}, \text{ so in every case we obtain a nonempty intersection.}$

3. Hyperrings and hyperfields

Definition 3.1 ([1], [2]). A hyperring is a tuple $(R, +, \cdot, 0, 1)$ which satisfies the following axioms:

- (R1) (R, +, 0) is a canonical hypergroup,
- (R2) $(R, \cdot, 1)$ is a commutaive monoid such that $x \cdot 0 = 0$ for all $x \in R$,
- (R3) the operation \cdot is distributive with respect to the hyperoperation +. That is, for all $x, y, z \in R$,

$$x(y+z) = xy + xz$$

as sets. Here for $x \in H$ and $A \subseteq H$ we have set

$$xA := \{xa \mid a \in A\}.$$

If $(R \setminus \{0\}, \cdot, 1)$ is an abelian group, then $(R, +, \cdot, 0, 1)$ is called hyperfield.

The following example is the original example of a quotient hyperring (see [8]).

EXAMPLE 3.2. Let R be a commutative ring with 1, G a normal subgroup of its multiplicative semigroup and recall the notations introduced in Example 2.7. One may define multiplication in R/G as $aG \cdot bG := (ab)G$. Then $(R/G, +, \cdot, \{0\}, 1_RG)$ is a hyperring and if R is a field, then it is a hyperfield. We have to show that (R3) holds. Take $a, b, c \in R$ and suppose that $xG \in aG \cdot (bG + cG)$. Then there exist $g, h \in G$ such that x = a(bg + ch) = abg + ach, where we have used the distributivity law in R. We obtain

$$xG \in \{(abg + ach)G \mid g, h \in G\} = abG + acG = aGbG + aGcG.$$

This shows that $aG \cdot (bG + cG) \subseteq aGbG + aGcG$. To show the converse inclusion we note that

$$xG \in aGbG + aGcG = (ab)G + (ac)G.$$

Hence there exist $g, h \in G$ such that x = (ab)g + (ac)h = a(bg + ch), where we used the distributivity law in R. We obtain

$$xG \in \{a(bg + ch)G \mid g, h \in G\} = \{aG(bg + ch)G \mid g, h \in G\} = aG \cdot (bG + cG).$$

If R is a field, then for every $a \in R$ we have that $a^{-1}G = (aG)^{-1}$. Indeed, by definition $aG \cdot a^{-1}G = (a \cdot a^{-1})G = 1_RG$.

EXAMPLE 3.3. As we have already noted in Remark 2.8 that the sign hypergroup H can be seen as a quotient $\mathbb{R}/\dot{\mathbb{R}}^2$. Since \mathbb{R} is a field, we obtain that $(H,+,\cdot,0,1)$ is a hyperfield, where we now denote * by + and \cdot behaves as follows:

$$-1 \cdot 1 = 1 \cdot -1 = -1,$$

 $0 \cdot 0 = 0 \cdot 1 = 1 \cdot 0 = 0 \cdot -1 = -1 \cdot 0 = 0,$
 $1 \cdot 1 = -1 \cdot -1 = 1.$

In [13] it is argued that the set X of order preserving mappings $f: H \to K$ such that the support of f is finite, where H is a ordered canonical hypergroup

and K an ordered hyperfield, is a hyperdomain, i.e., a hyperring without zero divisors, with the operations defined as follows:

$$(f+g)(x) := f(x) +_K g(x),$$

and

$$(fg)(x) := \sum_{x \in x_1 * x_2} f(x_1)g(x_2). \tag{3.1}$$

This seems to be an attempt to generalize the construction of formal power series. However, we first of all note that X is not even a hyperring since if f is non-constant and order preserving, then -f is not order preserving. In addition, if K is a hyperfield which is not a field, then the multiplication (fg)(x) defined in (3.1) is a subset of K and not an element of K as the definition of hyperring would require. Hence, we do not obtain a hyperring. Finally, even if we do not restrict to order preserving mappings and assume K to be a field, we show, in the next example, that the multiplication defined in (3.1) is not associative.

EXAMPLE 3.4. Let $H = \{-1,0,1\}$ denote the sign hypergroup and K an ordered field. We define the maps $f,g,h: H \to K$ as follows $f(-1) = f(0) = f(1) = 1_K$, $g(-1) = g(0) = g(1) = -1_K$, and $h(1) = 1_K$, $h(0) = -1_K$, $h(-1) = 0_K$. By direct computations we obtain

$$((fg)h)(1) = 5 \cdot (-1_K) + 5 \cdot 1_K + 3 \cdot (-1_K) + 5 \cdot (-1_K) = 8 \cdot (-1_K)$$

and

$$(f(gh))(1) = 2 \cdot (-1_K) + 2 \cdot (-1_K) + 2 \cdot (-1_K) = 6 \cdot (-1_K)$$

where we used the fact that

$$1 \in 1 * 1, 1 * 0, 0 * 1, 1 * -1, -1 * 1,$$

that

$$-1 \in -1 * -1, -1 * 0, 0 * -1, 1 * -1, -1 * 1$$

and that

$$0 \in 0 * 0, 1 * -1, -1 * 1.$$

Here * denotes the operation of H. We conclude that $f(gh) \neq (fg)h$, hence the associativity law does not hold.

4. Hypervaluations

As it was mentioned before, there are several approaches to the definition of a hypervaluation on a hyperfield. We now wish to investigate the following.

Definition 4.1 ([13]). Let $(F, +, \cdot, 0, 1)$ be a hyperfield and $(H, *, e, \leq)$ be an ordered canonical hypergroup. A surjective map $w : F \to H \cup \{\infty\}$ is called a hypervaluation on the hyperfield F if the following properties are satisfied:

(V1)
$$w(x) = \infty \iff x = 0$$
,

 $(V2) \ w(-x) = w(x),$

 $(V3) \ w(x \cdot y) \in w(x) * w(y),$

$$(V4) z \in x + y \implies w(z) \ge \min\{w(x), w(y)\}.$$

The following is a nontrivial example of a hypervaluation onto an ordered canonical hypergroup.

EXAMPLE 4.2. Let K be a field and Γ an ordered abelian group. Assume that a classical valuation

$$v: K \to \Gamma \cup \{\infty\}$$

is given. We now define a hypervaluation w from K onto the sign hypergroup $\{-1,0,1\}$.

$$w(x) = \begin{cases} 1 & \text{if } \infty \neq v(x) > 0_{\Gamma} \\ 0 & \text{if } v(x) = 0_{\Gamma} \\ -1 & \text{if } v(x) < 0_{\Gamma} \\ \infty & \text{otherwise} \end{cases}$$

Let us show that w is a hypervaluation on the field K, as in Definition 4.1. Clearly, $w(x) = \infty$ if and only if $x = 0_K$ and w(x) = w(-x) because that is true for v. In order to show that $w(xy) \in w(x) * w(y)$, where * denotes the operation in the sign hypergroup, we observe that if v(x) and v(y) have the same sign there is nothing to show, since v(xy) = v(x) + v(y) will have the same sign as v(x) and v(y). If v(x) = 0, then w(x) = 0 and v(xy) = v(y) and w(xy) = w(y) = 0 * w(y); the same holds if v(y) = 0. If x = 0 or y = 0, then the situation is clear. If, say, $v(x) < 0_{\Gamma}$ and $v(y) > 0_{\Gamma}$, then $w(xy) \in \{-1,0,1\} = -1 * 1 = w(x) * w(y)$, and similarly in the case where $v(x) > 0_{\Gamma}$ and $v(y) < 0_{\Gamma}$. This shows that the third axiom of a hypervaluation holds for w. The fourth and last axiom states:

$$z \in x + y \Longrightarrow w(z) \succeq \min\{w(x), w(y)\}\$$

where \leq denotes the ordering in the sign hypergroup. Since K is a field we have to check that $w(x+y) \succeq \min\{w(x), w(y)\}$. If $w(x) \neq w(y)$, then clearly $v(x) \neq v(y)$, so we have $v(x+y) = \min\{v(x), v(y)\}$. Hence $w(x+y) = \min\{w(x), w(y)\}$. If x=y=0 there is nothing to show. If w(x) = w(y) = 0, then $v(x) = v(y) = 0_{\Gamma}$, so $v(x+y) \geq 0_{\Gamma}$. Then $w(x+y) \in \{0,1,\infty\}$ and $w(x+y) \succeq 0 = \min\{w(x), w(y)\}$. If w(x) = w(y) = 1, then $v(x), v(y) > 0_{\Gamma}$, so $v(x+y) > 0_{\Gamma}$. We obtain $w(x+y) \in \{1,\infty\}$ and $w(x+y) \succeq 1 = \min\{w(x), w(y)\}$. Finally, if w(x) = w(y) = -1, then $w(x+y) \in \{-1, 0, 1, \infty\}$, but $\min\{w(x), w(y)\} = -1$, so the fourth axiom also holds for w.

Lemma 4.3. Let $w: F \to H \cup \{\infty\}$ be a hypervaluation. Then $w(1_F) = e$ and $w(x^{-1}) = (w(x))^{-1}$.

Proof. Let $x \in F$ be such that w(x) = e. Then $e = w(x \cdot 1_F) \in w(x) * w(1_F) = \{w(1_F)\}$, hence $e = w(1_F)$. To prove the second assertion observe that, since $xx^{-1} = 1_F$, we have that $w(xx^{-1}) = w(1_F) = e$. By axiom (V3) we obtain that $e = w(xx^{-1}) \in w(x) * w(x^{-1})$, then axiom (H3) implies that $w(x^{-1}) = (w(x))^{-1}$.

Definition 4.4. Let $(R, +, \cdot, 0, 1)$ be a hyperring. An element $x \in R$ is called a *unit of* R if there exists $y \in R$ such that $x \cdot y = 1$.

Definition 4.5. Let F be a hyperfield and $R \subseteq F$ a hyperring with respect to the hyperaddition and multiplication of F. If for every $x \in F$ either $x \in R$ or $x^{-1} \in R$, then R is called a *valuation hyperring*.

Definition 4.6 ([6], [2]). Let R be a hyperring.

- (1) A nonempty subset $I \subseteq R$ is a *hyperideal* if for all $a, b \in I$ and for all $r \in R$ we have $a + b \subseteq I$, $-a \in I$ and $ar \in I$.
- (2) A hyperideal $I \subsetneq R$ is maximal if I satisfies the following property: if $J \subseteq R$ is a hyperideal of R such that $I \subsetneq J$, then J = R.

In [13] one can find the first three statements of the following proposition. For the sake of completeness we rewrite the proof and complete it with details where needed. For our purposes, we also add a fourth statement.

Proposition 4.7. Let $(F, +, \cdot, 0, 1)$ be a hyperfield, $(H, *, e, \leq)$ an ordered canonical hypergroup and $w : F \to H \cup \{\infty\}$ a hypervaluation.

- 1) The set $\mathcal{O}_w = \{x \in F \mid w(x) \geq e\}$ is a valuation hyperring.
- 2) The set $U_w = \{x \in F \mid w(x) = e\}$ is a group under the multiplication of the hyperfield F and consists of all units of \mathcal{O}_w .
- 3) The set $\mathfrak{m}_w = \{x \in F \mid w(x) > e\}$ is the unique maximal hyperideal of \mathcal{O}_w .
- 4) The quotient $G := (F \setminus \{0\})/U_w$ is an ordered abelian group with respect to the operation $xU_w \cdot yU_w = xyU_w$ and the ordering $xU_w \leq yU_w \Leftrightarrow yx^{-1} \in \mathcal{O}_w$.

Proof. 1) If $x, y \in \mathcal{O}_w$, then $w(x), w(y) \geq e$, so by part 3) of Lemma 2.13 we obtain that for all $b \in w(x) * w(y)$ we have $b \geq e$. Since $w(xy) \in w(x) * w(y)$, in particular $w(xy) \geq e$, hence $xy \in \mathcal{O}_w$. The inclusion $x + y \subseteq \mathcal{O}_w$ follows from (V4) and from the fact that $\min\{w(x), w(y)\} \geq e$. Indeed by (V4),

$$z \in x + y \Longrightarrow w(z) \ge \min\{w(x), w(y)\} \ge e.$$

To see that \mathcal{O}_w is a valuation hyperring, take $x \in F \setminus \mathcal{O}_w$. Then w(x) < e, which implies that $w(x^{-1}) = (w(x))^{-1} > e$ by Lemma 4.3 and part 2) of Lemma 2.13. Thus, $x^{-1} \in \mathcal{O}_w$.

- 2) If $x \in U_w$, then $x^{-1} \in U_w$ too. Indeed, $w(x^{-1}) = (w(x))^{-1} = e$. Hence the elements of U_w are units of \mathcal{O}_w . To show the other inclusion, assume that $x \in \mathcal{O}_w \setminus U_w$. Then w(x) > e, so $w(x^{-1}) < e$ by part 2) of Lemma 2.13, which means that $x^{-1} \notin \mathcal{O}_w$.
 - 3) Let $x, y \in \mathfrak{m}_w$. Then by (V4) we obtain that

$$z \in x + y \Longrightarrow w(z) \ge \min\{w(x), w(y)\} > e.$$

Thus, $x+y \subseteq \mathfrak{m}_w$. From w(-x) = w(x) it follows that $-x \in \mathfrak{m}_w$. If $x \in \mathfrak{m}_w$ and $y \in \mathcal{O}_w$, then w(x) > e and $w(y) \ge e$, so by part 4) of Lemma 2.13 we obtain that for every $b \in w(x)*w(y)$ we have b > e. Since by (V3) $w(xy) \in w(x)*w(y)$, we conclude that $xy \in \mathfrak{m}_w$. We have proved that \mathfrak{m}_w is a hyperideal. It follows from part 2) and from the fact that if a hyperideal contains a unit, then is not proper, that \mathfrak{m}_w is the unique maximal hyperideal of \mathcal{O}_w .

4) By part 2) U_w is a subgroup of the abelian group $(F \setminus \{0\}, \cdot, 1)$. So G is the quotient group with respect to this (normal) subgroup. The proof that \leq is a linear order on G is exactly the same as in the classical case.

Definition 4.8 ([6]). Let $(H_1, *_1, e_1)$ and $(H_2, *_2, e_2)$ be canonical hypergroups.

- (1) A homomorphism of hypergroups is a function $f: H_1 \to H_2$ such that $f(e_1) = e_2$ and $f(a *_1 b) \subseteq f(a) *_2 f(b)$.
- (2) A strict homomorphism of hypergroups is a function $f: H_1 \to H_2$ such that $f(e_1) = e_2$ and $f(a *_1 b) = f(a) *_2 f(b)$ as sets.
- (3) An *isomorphism of hypergroups* is a strict homomorphism of hypergroups which is bijective.

In [13] it is claimed that two hypervaluations $v: F \to H_v \cup \{\infty\}$ and $w: F \to H_w \cup \{\infty\}$ induce the same valuation hyperring if and only if there exists an order preserving isomorphism f between the value hypergroups H_v, H_w such that $w = f \circ v$. Although this statement is true in classical valuation theory, this is no longer true for hypervaluations.

EXAMPLE 4.9. Consider the hypervaluations v, w from Example 4.2. We observe that the valuation ring \mathcal{O}_v of v coincides with the valuation (hyper)ring of w. Indeed, if $v(x) \geq 0_{\Gamma}$, then $w(x) \in \{0,1\}$, so $w(x) \succeq 0$ which means that $\mathcal{O}_v \subseteq \mathcal{O}_w$. On the other hand, if $w(x) \succeq 0$, then, by definition, $v(x) \geq 0$, so $x \in \mathcal{O}_w$ implies $x \in \mathcal{O}_v$. Note that there is no order preserving isomorphism between the ordered abelian group Γ and the sign hypergroup. But in the present case, as shown above, they define the same valuation (hyper)ring in K.

The next theorem states that any hypervaluation from a hyperfield onto an ordered canonical hypergroup is the composition of a hypervaluation onto an

ordered abelian group (which induces the same valuation hyperring) and an order preserving homomorphism of hypergroups.

Theorem 4.10. Let F be a hyperfield, $(H, *, e, \leq)$ an ordered canonical hypergroup and $w: F \to H \cup \{\infty\}$ a hypervaluation. Then there exists a hypervaluation $v: F \to G \cup \{\infty\}$, where G is an ordered abelian group, and an order preserving homomorphism $h: G \to H$ of hypergroups from G to H s.t. $w = h \circ v$ and $\mathcal{O}_v = \mathcal{O}_w$.

Proof. By part 4) of Proposition 4.7 we can consider the ordered abelian group $G = (F \setminus \{0\})/U_w$. Let $h: G \to H$ be the map

$$h(xU_w) = w(x).$$

We first show that h is well defined. Suppose that $xU_w = yU_w$. This is equivalent to $xy^{-1} \in U_w$, which holds if and only if $w(xy^{-1}) = e$. From (V3) and Lemma 4.3 we obtain that $e \in w(x) * w(y^{-1}) = w(x) * (w(y))^{-1}$ and this means that w(x) = w(y) by the uniqueness required in axiom (H3). This proves that h is well defined. To show that h is a homomorphism of hypergroups, note that $h(1U_w) = w(1) = e$ by Lemma 4.3. Take $xU_w, yU_w \in G$. According to Remark 2.5 we obtain that

$$h(xU_w \cdot yU_w) = \{h(xyU_w)\} = \{w(xy)\} \subseteq w(x) * w(y) = h(xU_w) * h(yU_w),$$

which proves that h is a homomorphism of hypergroups. Moreover, assume that $xU_w \leq yU_w$, which by definition is equivalent to $e \leq w(yx^{-1})$. From the latter it follows by condition (2.1) of Definition 2.10 that $\{w(x)\} = e * w(x) \nearrow w(yx^{-1}) * w(x)$. Hence by part 1) of Lemma 2.13, $w(x) \leq b$ for every $b \in w(yx^{-1}) * w(x)$. As $w(y) = w(yx^{-1}x) \in w(yx^{-1}) * w(x)$, we obtain that $w(x) \leq w(y)$. This proves that h is order preserving.

We now define $v: F \to G \cup \{\infty\}$ as follows:

$$v(x) = \begin{cases} xU_w, & \text{if } x \neq 0\\ \infty, & \text{if } x = 0. \end{cases}$$

The map v is a hypervaluation onto the ordered abelian group G (again, modulo the provision of Remark 2.5). The only axiom which needs justification is the fourth: if $z \in x + y$ for some $x, y \in F$, then we wish to show that $v(z) \ge \min\{v(x), v(y)\}$. Assume, without loss of generality, that $v(x) \le v(y)$, so that $yx^{-1} \in \mathcal{O}_w$. From $z \in x + y$ it follows that

$$yx^{-1} \in (x-z)x^{-1} = 1 - zx^{-1}.$$

This means that $zx^{-1} \in yx^{-1} - 1 \subseteq \mathcal{O}_w$, which proves that $v(z) \geq v(x)$.

Finally, one can see that $w = h \circ v$. Indeed, if $x \in F$, then if $x \neq 0$, we have that $h(v(x)) = h(xU_w) = w(x)$ by definition. If x = 0, the situation is clear once we set $h(\infty) := \infty$. It is left to show that $\mathcal{O}_v = \mathcal{O}_w$. We observe that

 $x \in \mathcal{O}_w$ if and only if $1U_w \le xU_w$, which happens if and only if $x \in \mathcal{O}_v$. This completes the proof.

Acknowledgments

The authors would like to thank prof. F.-V. Kuhlmann and K. Kuhlmann for their careful reading and their many extremely useful remarks which helped to improve this paper significantly.

References

- P. Corsini, V. Leoreanu-Fotea, Applications of Hyperstructure Theory, Advances in Mathematics, 5, Kluwer Academic Publishers, Dordrecht, 2003.
- B. Davvaz, V. Leoreanu-Fotea, Hyperring Theory and Applications, International Academic Press, Palm Harbor, USA, 2007.
- S. Omidi, B. Davvaz, Ordered Krasner Hyperrings, Iranian Journal of Mathematical Sciences and Informatics, 12(2), (2017), 35-49.
- B. Davvaz, A. Salasi, A Realization of Hyperrings, Communications in Algebra, 34(12), (2006), 4389-4400.
- J. Flenner, Relative Decidability and Definability in Henselian Valued Fields, Journal of Symbolic Logic, 76(4), (2011), 1240-1260.
- J. Jun, Algebraic Geometry Over Hyperrings, Advances in Mathematics, 323, (2018), 142-192.
- M. Krasner, Approximation Des Corps Valués Complets de Caractéristique p ≠ 0 Par Ceux de Caractéristique 0, Colloque d'Algèbre supérieure, Bruxelles, 1956, pp. 129-206,.
- M. Krasner, A Class of Hyperrings and Hyperfields, International Journal of Mathematics and Mathematical Sciences, 6, (1983), 307-311.
- F.-V. Kuhlmann, Quantifier Elimination for Henselian Fields Relative to Additive and Multiplicative Congruences, Israel Journal of Mathematics, 85, (1994), 277-306.
- J. Lee, Hyperfields, Truncated DVRs and Valued Fields, Journal of Number Theory, 212, (2020), 40-71.
- M. Marshall, Real Reduced Multirings and Multifields, Journal of Pure and Applied Algebra, 205, (2006), 452-468.
- 12. F. Marty, Sur Une généralisation de La Notion de Groupe, 8th Congress Math. Scandenaves, Stockholm, 1934, pp. 45-49.
- Kh. Mirdar Harijani, S. M. Anvariyeh, A Hypervaluation of a Hyperfield Onto a Totally Ordered Canonical Hypergroup, Studia Scientiarum Mathematicarum Hungarica, 52(1), (2015), 87-101.