

On the Notion of Fuzzy Shadowing Property

Mehdi Fatehi Nia

Department of Mathematics, Yazd University, 89195-741 Yazd, Iran.

E-mail: `fatehiniam@yazd.ac.ir`

ABSTRACT. This paper is concerned with the study of fuzzy dynamical systems. Let $(X, M, *)$ be a fuzzy metric space in the sense of George and Veeramani. A fuzzy discrete dynamical system is given by any fuzzy continuous self-map defined on X . We introduce the various fuzzy shadowing and fuzzy topological transitivity on a fuzzy discrete dynamical systems. Some relations between these notions have been proved.

Keywords: Fuzzy metric, Fuzzy discrete dynamical systems, Fuzzy shadowing, Fuzzy ergodic shadowing, Fuzzy topological mixing.

2000 Mathematics subject classification: 37C50, 37C15.

1. INTRODUCTION

Fuzzy topology is an extension of the ordinary topology which has become an area of active research in the recent years, because of its numerous applications [2, 21]. George and Veeramani (see [8, 9]) modified and studied a notion of fuzzy metric M on a set X via of continuous t -norms which introduced by Kramosil and Michalek [15]. From now on, when we talk about fuzzy metrics we refer to this type of fuzzy metric spaces.

George and Veeramani proved that M induces a topology on X . This topology is not the same as the fuzzy topology. Actually, this topology also can be constructed on each fuzzy metric space in the sense of Kramosil and Michalek [12, 15, 18]. We should remark that the topology on X deduced from a fuzzy metric M , does not depend on the t -norm [9].

Received 29 May 2015; Accepted 28 June 2016

©2018 Academic Center for Education, Culture and Research TMU

The shadowing property and chain mixing are two of the most important concepts in discrete dynamical systems [1], which are closely related to stability and chaotic behavior of dynamical systems, see, for instance [14, 24, 25, 26, 27] and they are essential parts of stability and ergodic theory [1]. In this direction the concept of ergodic shadowing property, topological mixing and chain mixing are another formal ways to get some special dynamical properties, for discrete dynamical system (X, f) , where f is an onto continuous map (see [3, 4, 19, 20, 23, 25, 26]).

The study of discrete fuzzy dynamical systems has been done by many authors and different properties of fuzzy discrete dynamical systems were considered. For example, study of some chaotic properties of the discrete fuzzy dynamical system has been done by some authors [2, 16, 17, 22].

Nearness depends on the time and we have not introduced any distance as a metric. In fact, fuzzy metric determines the rate of nearness without introducing a distance number, this is the main idea of the fuzzy metric spaces. This theory implies that *the rate of nearness is more important than the notion of distance*. For this reason we use of the word fuzzy, since there is not any distance in fuzzy metric space theory. In this paper we have extend the notion of shadowing by using of the rate of nearness instead of distance. In coding theory distance is not reachable, but we can determine the rate of nearness. Since shadowing is one of the important property in the theory of discrete dynamical systems as it is close to the stability of the system and also to the chaotic behavior of the systems [6, 7, 14, 25, 27], this extension can be a useful tool to study stability and chaos in fuzzy discrete dynamical systems.

The present paper is organized as follows: In Section 2, several basic definitions and notations are presented. In Section 3, we consider the concept of fuzzy discrete dynamical systems which is a generalization of discrete dynamical systems [2, 17]. We use F -.... instead of Fuzzy....

The notions of δ - F -pseudo-orbit, F -shadowing property, F -topological mixing, F -chain transitive, F -ergodic shadowing property, have been developed and presented. Some propositions on the above notions have also been established. Actually, we show that the following statements hold:

- 1) Any mapping with F -ergodic shadowing property is F -chain transitive.
- 2) If f has the F -ergodic shadowing property then for any natural number k the mapping f^k has also the F -ergodic shadowing property.
- 3) F -ergodic shadowing property implies F -shadowing property.
- 4) If f has the F -shadowing and F -chain mixing properties then it has the F -topologically mixing property.

In Section 4 we have introduced few examples to illustrate the definitions above. A tent map is one of the most popular and the simplest chaotic maps [4]. In Example 4.1 and Remark 4.2, we consider the family of tent maps and show that they have the

F -shadowing and F -chain mixing properties. Also, in Example 4.3 we show that the classical shadowing property is a special case of fuzzy shadowing property and they are essentially different.

As usual the paper finishes some concluding remarks and open problems (Section 5).

2. PRELIMINARIES

This section contains two major subsections: Fuzzy Metric Space and Discrete Dynamical Systems. We start this section by recalling some pertinent concepts.

Fuzzy Metric Space

Let $*$ be a continuous t -norm, i.e., it is a binary operation from $[0, 1] \times [0, 1]$ to $[0, 1]$ with the following conditions:

- i) $*$ is associative and commutative;
- ii) $*$ is continuous;
- iii) $a * 1 = a$ for all $a \in [0, 1]$;
- iv) $a * b \leq c * d$ whenever $a \leq c$, $b \leq d$ and $a, b, c, d \in [0, 1]$.

Lemma 2.1. [8]. *For any $r_1 > r_2$, we can find r_3 such that $r_1 * r_3 \geq r_2$ and for any r_4 we can find r_5 such that $r_5 * r_5 \geq r_4$, ($r_1, r_2, r_3, r_4, r_5 \in (0, 1)$).*

We also assume that $(X, M, *)$ is a *fuzzy metric space* [8, 9] i.e., X is a nonempty set, $*$ is a continuous t -norm and

$M : X \times X \times (0, \infty) \rightarrow [0, 1]$ is a mapping with the following properties:

For every $x, y, z \in X$ and $t, s > 0$

- 1) $M(x, y, t) > 0$;
- 2) $M(x, y, t) = 1$ if and only if $x = y$;
- 3) $M(x, y, t) = M(y, x, t)$;
- 4) $M(x, z, t + s) \geq M(x, y, t) * M(y, z, s)$;
- 5) $M(x, y, \cdot) : (0, \infty) \rightarrow [0, 1]$ is a continuous map.

If $(X, M, *)$ is a fuzzy metric space, we will say that M is a fuzzy metric on X . The function $M(x, y, t)$ denotes the degree of nearness between x and y respect to t . Many interesting examples of fuzzy metric spaces can be found in [11].

Lemma 2.2. [10]. *$M(x, y, \cdot) : (0, \infty) \rightarrow (0, 1]$ is nondecreasing for all x, y in X .*

Definition 2.3. [8] Let $(X, M, *)$ be a fuzzy metric space. A set $A \subset X$ is called a *fuzzy open set* if for any $x \in A$ there exists $0 < r < 1$ and $T_0 \in (0, \infty)$ such that if $M(x, y, t) > 1 - r$, for all $t > T_0$ then $y \in A$.

Since $M(x, y, t) \geq M(x, y, T_0)$ for all $t > T_0$, then in the above definition it is sufficient that for one $T_0 \in (0, \infty)$ if $M(x, y, T_0) > 1 - r$ then $y \in A$.

Definition 2.4. [9]. Let $(X, M, *)$ be a fuzzy metric space. For $t > 0$, the open ball $B(x, r, t)$ with center $x \in X$ and radius $0 < r < 1$ is defined by:

$$B(x, r, t) = \{y \in X; M(x, y, t) > 1 - r\}.$$

Similarly, for $t > 0$, the closed ball with the center $x \in X$ and radius $0 < r < 1$ is defined by:

$$B[x, r, t] = \{y \in X; M(x, y, t) \geq 1 - r\}.$$

In [10] it has been proved that every fuzzy metric M on X generates a Hausdorff first countable topology τ_M on X which has a base the family of open sets of the form

$$\{B(x, r, t) : x \in X, r \in (0, 1), t > 0\}.$$

Functions $M(x, y, t)$ presented in Examples from 2.5 to 2.7 will be utilized in the Section 4.

EXAMPLE 2.5. [9] Let (X, d) be a metric space. Denote by $a.b$ the usual multiplication for all $a, b \in [0, 1]$, and let M_d be the function defined on $X \times X \times (0, \infty)$ by $M_d(x, y, t) = \frac{t}{t + d(x, y)}$. Then $(X, M_d, .)$ is a fuzzy metric space called standard fuzzy metric space, and $(M_d, .)$ will be called the standard fuzzy metric of d . And the topology τ_{M_d} generated by d coincides with the topology τ_{M_d} generated by the fuzzy metric M_d .

EXAMPLE 2.6. [11] Let $X = (0, 1]$ and let $\varphi : \mathbb{R}^+ \rightarrow (0, 1]$ be a function given by $\varphi(t) = t$ if $t \leq 1$ and $\varphi(t) = 1$ elsewhere. Define the function M on $X \times X \times (0, \infty)$ by:

$$M(x, y, t) = \begin{cases} 1 & \text{if } x = y, \\ \frac{\min\{x, y\}}{\max\{x, y\}} \cdot \varphi(t) & \text{if } x \neq y. \end{cases}$$

It is easy to verify that $(X, M, .)$ is a fuzzy metric on X , where $a.b = ab$ for all $a, b \in X$.

Notice that $B(x, \frac{1}{2}, \frac{1}{2}) = x$ for each $x \in X$ and so τ_M is the discrete topology.

EXAMPLE 2.7. [11] Let $X = (0, 1]$ and define the function M_1 on $X \times X \times (0, \infty)$ by:

$$M_1(x, y, t) = \begin{cases} 1 & \text{if } x = y, \\ \frac{\min\{x, y\}}{\max\{x, y\}} & \text{if } x \neq y. \end{cases}$$

It is easy to verify that $(X, M_1, .)$ is a fuzzy metric on X , where $a.b = ab$ for all $a, b \in X$. Let $\epsilon > 0$ and $a \in X$. This is clear that $B(a, \epsilon, t) \subset B_\epsilon(a)$ and $B_\epsilon(a\epsilon) \subset B(a, \epsilon, t)$, for all $t \in \mathbb{R}^+$. So τ_{M_1} is the usual topology of \mathbb{R} .

Theorem 2.8. [9]. Let $(X, M, *)$ be a fuzzy metric space and τ_M be the topology induced by the fuzzy metric. Then for a sequence $\{x_n\}_{n=1}^{\infty}$ in X , $\lim_{n \rightarrow \infty} x_n = x$ if and only if $\lim_{n \rightarrow \infty} M(x_n, x, t) = 1$ for all $t > 0$.

A sequence $\{x_n\}_{n=1}^{\infty}$ in a fuzzy metric space $(X, M, *)$ is a Cauchy sequence if and only if for each $\epsilon, t > 0$ there exists $n_0 \in \mathbb{N}$ such that $M(x_n, x_m, t) > 1 - \epsilon$ for all $n, m \geq n_0$. A fuzzy metric space in which every Cauchy sequence is convergent is called a complete fuzzy metric space.

Definition 2.9. [8] Let $(X, M, *)$ be a fuzzy metric space and $f : X \rightarrow X$ be a fuzzy map. Then f is said to be a *fuzzy continuous* map at the point x_0 , if for any $\epsilon \in (0, 1)$ and $t > 0$ there exists $\delta \in (0, 1)$ and $t' > 0$ such that for each x with $M(x, x_0, t') > 1 - \delta$. So $M(f(x), f(x_0), t) > 1 - \epsilon$.

f is said to be a fuzzy continuous map if f is fuzzy continuous at any point of X .

Discrete dynamical systems. Here, we recall some notions and results of discrete dynamical systems which we require in this paper [4, 5].

Let (X, d) be a compact metric space and $f : X \rightarrow X$ be a continuous map. For any two open subsets U and V of X , denote $N(U, V, f) = \{m \in \mathbb{N}; f^m(U) \cap V \neq \emptyset\}$. When there is no ambiguity, we denote it by $N(U, V)$. We say that f is *topologically transitive* if for every pair of open sets U and V in X , $N(U, V) \neq \emptyset$. *Topological mixing* means that for any two open subsets U and V , the set $N(U, V)$ contains any natural number $n \geq n_0$, for some fixed $n_0 \in \mathbb{N}$.

Given a number $\delta > 0$, a δ -pseudo orbit for f is a sequence $\{x_i\}_{1 \leq i \leq b}$ such that $d(f(x_i), x_{i+1}) < \delta$, for every $1 \leq i \leq b$. If $b < \infty$, then we say that the finite δ -pseudo orbit $\{x_i\}_{1 \leq i \leq b}$ of f is a δ -chain of f from x_1 to x_b of length b . A point $x \in X$ is called a *chain recurrent point* of f if for every $\delta > 0$, there is a δ -chain from x to x . Denote by $CR(f)$ the set of all chain recurrent points of f . A sequence $\{x_i\}_{1 \leq i \leq b}$ is said to be ϵ -shadowed by a point x in X if $d(f^i(x), x_i) < \epsilon$ for each $1 \leq i \leq b$. A mapping f is said to have *shadowing property* if for any $\delta > 0$, there is a $\epsilon > 0$ such that every δ -pseudo orbit of f can be ϵ -shadowed by some point in X .

A mapping f is called *chain transitive* if for any two points $x, y \in X$ and any $\delta > 0$ there exists a δ -chain from x to y . The mapping f is called chain mixing if for any two points $x, y \in X$ and any $\delta > 0$, there is a positive integer n_0 such that for any integer $n \geq n_0$ there is a δ -chain from x to y of length n .

Given a sequence $\eta = \{x_i\}_{i \in \mathbb{N}}$, denote

$$Npo(\eta, \delta) = \{i | d(f(x_i), x_{i+1}) \geq \delta\}$$

and

$$Npo_n(\eta, \delta) = Npo(\eta, \delta) \cap \{1, \dots, n-1\}.$$

For a sequence η and a point x of X , denote

$$Ns(\eta, x, \delta) = \{i | d(f^i(x), x_i) \geq \delta\}$$

and

$$Ns_n(\eta, x, \delta) = Ns(\eta, x, \delta) \cap \{1, \dots, n-1\}.$$

Definition 2.10. [5]. A sequence η is a δ -ergodic pseudo orbit if $Npo(\eta, \delta)$ has density zero, it means that $\lim_{n \rightarrow \infty} \frac{\text{card}(Npo_n(\eta, \delta))}{n} = 0$, where $\text{card}(\cdot)$ denotes the cardinal number. A δ -ergodic pseudo orbit η is said to be ϵ -ergodic shadowed by a point x in X if $\lim_{n \rightarrow \infty} \frac{\text{card}(Ns_n(\eta, x, \epsilon))}{n} = 0$.

In other words, $Ns(\eta, x, \epsilon)$, has density zero. A mapping f has ergodic shadowing property if for any $\epsilon > 0$ there is a $\delta > 0$ such that any δ -ergodic pseudo orbit of f can be ϵ -ergodic shadowed by some point in X .

The shadowing property and chain recurrence is one of the most important concepts in dynamical systems [1], which are closely related to stability and chaos of systems, see, for instance [14, 25, 26] and are essential parts of stability and ergodic theory [1].

In [5] Fakhari and Ghane prove the following theorem:

Theorem 2.11. [5] *Let f be a continuous onto map of a compact metric space X . For the dynamical system (X, f) , the following properties are equivalent:*

- (a) *ergodic shadowing,*
- (b) *shadowing and chain mixing,*
- (c) *shadowing and topological mixing. We are going to study other concepts and their relations in another papers.*

3. FUZZY DYNAMICAL SYSTEMS

We begin this section by some definitions which is an extension to fuzzy topology of the concepts in the previous section and will be used in Theorems 3.4 to 3.7 and Section 4.

Suppose X is a nonempty set and $f : X \rightarrow X$ is an arbitrary map. The n -th iteration f^n of the map f is defined inductively by $f^1 = f$ and $f^n = f \circ f^{n-1}$ for $n \geq 2$ and the orbit of x under f is the set of points $\{x, f(x), f^2(x), \dots\}$. It is well known [17] that (X, f) induces a dynamical system $(\mathbb{F}(X), \hat{f})$, called fuzzy discrete dynamical system, that is defined on the space $\mathbb{F}(X)$ of all fuzzy compact subsets of X . The map \hat{f} is usually called fuzzification (see [2] and [17] for more details. Now, we consider a fuzzy discrete dynamical system, denoted by $(X, M, *, f)$, as a fuzzy continuous map $f : X \rightarrow X$ with all iterations and identity map, where $(X, M, *)$ is a fuzzy metric space.

Definition 3.1. Given a number $\delta > 0$. A sequence $\{x_i\}_{i=0}^b$ of points in X is called a $\delta - F$ -pseudo-orbit of f if there exists

$T_0 \in (0, \infty)$ such that $M(f(x_i), x_{i+1}, T_0) > 1 - \delta$ for all $0 \leq i < b$.

If $b < \infty$, then we say that the finite $\delta - F$ -pseudo-orbit $\{x_i\}_{i=0}^b$ of f is a $\delta - F$ -chain of f from x_0 to x_b .

We say that a fuzzy dynamical system $(X, M, *, f)$ has the

F -shadowing property, if for any $\epsilon > 0$ there is a $\delta > 0$ such that every $\delta - F$ -pseudo-orbit $\{x_i\}_{i=0}^b$ of f can be $\epsilon - F$ -traced by some point in X that is:

There is $T_0 > 0$ and $x \in X$ such that $M(f^i(x), x_i, T_0) > 1 - \epsilon$, for each $0 \leq i \leq b$.

Let f be a fuzzy continuous map. For any two fuzzy open subsets U and V of X , let

$$N^f(U, V) = \{m \in \mathbb{N}; f^m(U) \cap V \neq \emptyset\}.$$

We say that f is F -topological transitive if for any two fuzzy open subsets U and V of X , $N^f(U, V) \neq \emptyset$. F -Topological mixing means that for any two fuzzy open subsets U and V , the set $N^f(U, V)$ contains any natural number $n \geq n_0$, for some fixed $n_0 \in \mathbb{N}$. From here the section we assume that $(X, M, *)$ is a complete and compact fuzzy metric space, $\mathbb{R}^+ = (0, \infty)$ and \mathbb{N} the positive integers. For brevity, we write the fuzzy metric spaces as X , wherever there is no risk of confusion.

Definition 3.2. A mapping f is called F -chain transitive if for any two points $x, y \in X$ and any $\delta > 0$ there exists a $\delta - F$ -chain from x to y . The mapping f is called F -chain mixing if for any two points $x, y \in X$ and any $\delta > 0$, there is a positive integer N_0 such that for any integer $n \geq N_0$ there is a $\delta - F$ -chain from x to y of length n .

Given a sequence $\mu = \{x_i\}_{i=0}^\infty$, let

$$N^f(\mu, \delta, t) = \{i; M(f(x_i), x_{i+1}, t) \leq 1 - \delta\}$$

and

$$N_n^f(\mu, \delta, t) = N^f(\mu, \delta, t) \cap \{1, \dots, n-1\}.$$

For a sequence μ and a point x of X , let

$$N_s^f(\mu, x, \delta, t) = \{i; M(f^i(x), x_i, t) \leq 1 - \delta\}$$

and

$$N_{sn}^f(\mu, x, \delta, t) = N_s^f(\mu, x, \delta, t) \cap \{1, \dots, n-1\}.$$

Definition 3.3. A sequence μ is a $\delta - F$ -ergodic pseudo orbit, if there exist $T_0 \in (0, \infty)$ such that

$$\lim_{n \rightarrow \infty} \frac{\text{card}(N_n^f(\mu, \delta, T_0))}{n} = 0.$$

A $\delta - F$ -ergodic pseudo orbit, is said to be $\epsilon - F$ -ergodic shadowed by a point x in X if

$$\lim_{n \rightarrow \infty} \frac{\text{card}(N_{sn}^f(\mu, x, \delta, T_0))}{n} = 0.$$

A mapping f has F -ergodic shadowing property if for any $\epsilon > 0$ there is a $\delta > 0$ such that any $\delta - F$ -ergodic pseudo orbit of f can be $\epsilon - F$ -ergodic shadowed by some point in X .

Now we are ready to show the results of this paper.

Theorem 3.4. *Any mapping with F -ergodic shadowing property is F -chain transitive.*

Proof. Let f be a mapping that has the F -ergodic shadowing property and $x, y \in X$. Suppose ϵ is an arbitrary positive number and $\delta > 0$ is a number corresponding to ϵ by the F -ergodic shadowing property.

Let $\{[a_i, b_i]; i = 0, 1, 2, \dots\}$ be a family of intervals such that $a_0 = 0, b_0 = 1$ and $a_k = b_{k-1} + k, b_k = a_k + k + 1$ for all $k \geq 1$. This is clear that the set $\{a_i, b_i; i = 0, 1, 2, \dots\}$ has density zero in natural numbers. Define the sequence $\{x_i\}_{i=0}^\infty$ by the following formulas:

$$x_i = \begin{cases} f^{i-a_k}(x) & \text{if } a_k \leq i < b_k, \\ f^{i-b_k}(x_k) & \text{if } b_k \leq i < a_{k+1}. \end{cases}$$

By definition of $\{x_i\}_{i=0}^\infty$, we have $M(f(x_i), x_{i+1}, t) = 1$ where $i = a_k$ or $i = b_k$ for some $k \geq 0$. This implies that $N^f(\mu, \delta, t)$ is a subset of $\{a_k, b_k; k = 0, 1, 2, \dots\}$. Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{N^f(\mu, \delta, t)}{n} &\leq \lim_{n \rightarrow \infty} \frac{\text{card}\{a_k, b_k; k = 0, 1, 2, \dots\} \cap \{0, 1, 2, \dots, n-1\}}{n} = 0. \end{aligned}$$

Where $\text{card}\{a_k, b_k; k = 0, 1, 2, \dots\}$ is cardinality of

$\{a_k, b_k; k = 0, 1, 2, \dots\}$. Then $\mu = \{x_i\}_{i=0}^\infty$ is a $\delta - F$ -ergodic pseudo orbit and can be $\epsilon - F$ -ergodic shadowed by some point z in X . Since $N_s^f(\mu, z, \epsilon, t)$ has density zero, for some $t \in (0, \infty)$. Then we can find $i, j, l, s \in \mathbb{N}$ with $j < l$ such that $M(f^j(x), f^i(z), t) \leq 1 - \epsilon, M(f^l(z), w, t) \leq 1 - \epsilon$ and $w \in f^{-s}(y)$.

Therefore

$$\{x, f(x), \dots, f^{j-1}(x), f^i(z), \dots, f^{l-1}(z), w, f(w), \dots, f^{s-1}(w), y\}$$

is an $\epsilon - F$ -pseudo orbit from x to y . □

Theorem 3.5. *If f has the F -ergodic shadowing property then for any natural number k the mapping f^k has also the F -ergodic shadowing property.*

Proof. Let f has the F -ergodic shadowing property. Given $\epsilon > 0$, let $\delta > 0$ be an ϵ modulus of F -ergodic shadowing property of f . Suppose $\{x_i\}_{i=0}^\infty$ is an $\delta - F$ -ergodic pseudo orbit for f^k . So there exist $T_0 \in (0, \infty)$ such that

$\lim_{n \rightarrow \infty} \frac{|N_n^{f^k}(\{x_i\}_{i=0}^\infty, x, \delta, T_0)|}{n} = 0$. We define the sequence $\{z_j\}_{j=0}^\infty$ with $z_{ki+l} = f^l(x_i)$ for $0 \leq l < k, n \in \mathbb{Z}_+$, that is

$$\{z_j\}_{j=0}^\infty = \{x_0, f(x_0), \dots, f^{k-1}(x_0), x_1, f(x_1), \dots, f^{k-1}(x_1), \dots\}.$$

It is easy to see that:

$$M(f(z_{ki+l}), z_{ki+l+1}, T_0) = \begin{cases} 0 & \text{if } 0 \leq l < k-1, \\ M(f^k(x_i), x_{i+1}, T_0) & \text{if } l = k-1. \end{cases}$$

So $\{z_j\}_{j=0}^\infty$ is an $\delta - F$ -ergodic pseudo orbit, f . Then there exists $x \in X$ such that $\{z_j\}_{j=0}^\infty$ is $\epsilon - F$ -shadowed by x in dynamical system $(X, M, *, f)$. Since $M(f^{ki}(x), z_{ki}, T_0) = M((f^k)^i(x), x_i, T_0)$, so

$$\text{card}(N_{sn}^{f^k}(\{x_i\}_{i=0}^\infty, x, \delta, T_0)) \leq \text{card}(N_{skn}^f(\{z_j\}_{j=0}^\infty, x, \delta, T_0)).$$

This implies that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\text{card}(N_{sn}^{f^k}(\{x_i\}_{i=0}^\infty, x, \delta, T_0))}{n} \\ \leq k \lim_{n \rightarrow \infty} \frac{\text{card}(N_{skn}^f(\{z_j\}_{j=0}^\infty, x, \delta, T_0))}{kn} = 0. \end{aligned}$$

□

Theorem 3.6. *Let f be a F -continuous onto map. For the fuzzy dynamical system $(X, M, *, f)$, if f has the F -ergodic shadowing property then it has the F -shadowing property.*

Proof. First, we prove that any finite F -pseudo orbit of f can be shadowed by true orbit. Let $\mu = \{z_j\}_{j=0}^n$ be a finite $\delta - F$ -pseudo orbit, by Theorem 3.5, we can find a $\delta - F$ -pseudo orbit

$\lambda = \{z_n = y_0, \dots, y_m = z_0\}$ from z_n to z_0 . Then

$\omega = \{z_0, \dots, z_n, y_1, \dots, y_{m-1}, z_0, z_1, \dots, z_n, y_1, \dots\}$ is a $\delta - F$ -ergodic pseudo orbit and then can be F -ergodic shadowed by some point $x \in X$. If the set $N_s^f(\omega, x, \epsilon, T_0)$ meet every μ interval then it would have positive density that is a contradict. Hence there exist at least one $k \in \mathbb{N}$ such that $M(f^{k+i}(x), z_i, t) > 1 - \epsilon$, for every $0 \leq i \leq n$. In other words μ is $\epsilon - F$ -shadowed by $f^k(x)$.

Now suppose $\{z_j\}_{j=0}^\infty$ is an $\delta - F$ -pseudo orbit of f , for $k > 0$, denote $x_i = z_i$ ($0 \leq i \leq k$). Then $\{x_i\}_{i=0}^k$ is an $\delta - F$ -pseudo orbit of f , and so is $\epsilon - F$ -shadowed by some point $w_k \in X$. Let $z \in X$ be an accumulation point of $\{w_k\}_{k=1}^\infty$, because X is a compact and complete fuzzy metric space. We show that $\{z_j\}_{j=0}^\infty$ is $\epsilon - F$ -shadowed by z .

Fix $n \geq 0$ and $t > 0$. By Lemma 2.1, there exist $\epsilon' > 0$ such that $(1 - \epsilon')(1 - \epsilon') \geq 1 - \epsilon$. Since f^n is a fuzzy continuous map then there exist $\delta' > 0$ and $t' > 0$ such that if $M(z, x, t') > 1 - \delta'$ then $M(f^n(z), f^n(x), t) > 1 - \epsilon'$, for every $x \in X$.

Take sufficiently large k_n satisfying $k_n > n$ and $M(z, w_{k_n}, t') > 1 - \delta'$. Then

$$\begin{aligned} M(f^n(z), z_n, 2t) &\geq M(f^n(z), f^n(w_{k_n}), t) * M(f^n(w_{k_n}), z_n, t) \\ &\geq (1 - \epsilon') * (1 - \epsilon') \\ &\geq 1 - \epsilon. \end{aligned}$$

This prove that $\{z_j\}_{j=0}^{\infty}$ is 2ϵ -shadowed by z . \square

Theorem 3.7. *If f has the F -shadowing and F -chain mixing properties then it has F -topologically mixing property.*

Proof. Suppose that f has the F -chain mixing property. Given two fuzzy open subsets U and V of X . Choose $x \in U$, $y \in V$, $t > 0$ and $\epsilon > 0$ such that $B(x, \epsilon, t) \subset U$ and $B(y, \epsilon, t) \subset V$. Let $\delta > 0$ be an ϵ -modulus of F -shadowing property. Since f is F -chain mixing then there is a positive integer N_0 such that for any integer $n \geq N_0$, there is a δ - F -chain, $x = x_1, x_2, \dots, x_n = y$, of length n from x to y . By F -shadowing property we can find $z \in X$ such that $M(f^i(z), x_i, t) > 1 - \epsilon$, for all $1 \leq i \leq n$. Particulary $M(z, x, t) > 1 - \epsilon$ and $M(f^n(z), y, t) > 1 - \epsilon$. So $z \in U$ and $f^n(z) \in V$. Then $f^n(U) \cap V \neq \emptyset$. \square

4. EXAMPLES

In the following, we give an interval map that has the F -topological mixing property.

EXAMPLE 4.1. *Let $X = [0, 1]$ and $a * b = ab$ for all $a, b \in X$. Let $(X, M_d, *)$ be the standard fuzzy metric space induced by d , where $d(x, y) = |x - y|$ for all $x, y \in X$. Suppose $f : X \rightarrow X$ is the tent map which is defined by:*

$$f(x) = \begin{cases} 2x & \text{if } 0 \leq x \leq \frac{1}{2}, \\ 2(1-x) & \text{if } \frac{1}{2} \leq x \leq 1. \end{cases}$$

It is known that the tent map has the shadowing and chain mixing properties (see [4, 20]) and we show that f has the F -shadowing and F -chain mixing properties.

Let $\epsilon > 0$. It is clear that there is $T_0 > 0$, such that $M_d(x, y, T_0) > 1 - \epsilon$ for all $x, y \in X$. So f has the F -shadowing and F -chain-mixing properties. It follows from Theorem 3.7 that f has the F -topological mixing property.

Remark 4.2. [4] Consider the family of tent maps, i.e., the piecewise linear maps $f_\beta(x) : [0, 1] \rightarrow [0, 1]$, $\sqrt{2} \leq \beta \leq 2$, defined by:

$$f_\beta(x) = \begin{cases} \beta x & \text{if } 0 \leq x \leq \frac{1}{2}, \\ \beta(1-x) & \text{if } \frac{1}{2} \leq x \leq 1. \end{cases}$$

In [4], the authors proved that, tent maps have the shadowing property for almost all parameters β , although they fail to have the shadowing property for an uncountable, dense set of parameters. But, similar to Example 4.1, for an arbitrary positive number ϵ , $M_d(x, y, T_0) > 1 - \epsilon$ for all $x, y \in X$. So $(X, M_d, *, f_\beta)$ has the F -shadowing property, for all $\sqrt{2} \leq \beta \leq 2$.

From Example 4.1 it can be seen that for metric M_d any continuous map has the F -shadowing property. Since the topology τ_d generated by d coincides with the topology τ_{M_d} generated by the fuzzy metric M_d , so the metric M_d is not a help for studying the dynamics.

The following example shows that the F -shadowing property is not completely similar to the shadowing property, although fuzzy metric space $(X, M, *)$ define the usual topological structure on X .

EXAMPLE 4.3. Let $X = (0, 1]$ and $f : X \rightarrow X$ be the following function (see Figure 1):

$$f(x) = \begin{cases} \frac{3}{4}x + \frac{1}{8} & \text{if } 0 < x \leq \frac{1}{2}, \\ \frac{3}{2}x - \frac{1}{4} & \text{if } \frac{1}{2} \leq x \leq \frac{3}{4}, \\ \frac{1}{2}x + \frac{1}{2} & \text{if } \frac{3}{4} \leq x \leq 1. \end{cases}$$

This kind of discrete dynamical systems is an important example to investigate various shadowing properties; see [8 Example 5.4.] and [11, Example 4.2.] for more details.

(a) It is well known that f does not have the shadowing property [3].

(b) Suppose $(X, M_d, *)$ is the standard fuzzy metric space in Example 2.5. It is clear that f is a fuzzy continuous function, respect to the fuzzy metric M_d .

Since $|x - y| < 1$ for all $x, y \in X$. Thus, given $\epsilon > 0$ there exist $T_0 > 0$ such that $M_d(x, y, t) > 1 - \epsilon$ for all $x, y \in X$ and $t > T_0$. So this is clear that for every $\epsilon - F$ -pseudo orbit, $\{x_i\}_{i=0}^\infty$, $M_d(f^i(\frac{1}{2}), x_i, t) > 1 - \epsilon$ for all $i \geq 0$ and $t > T_0$. Therefore $(X, M_d, *, f)$ has the F -shadowing property.


(c) Now we show that $(X, M, *, f)$ does not have the F -shadowing property, where $(X, M, *)$ is fuzzy metric space in Example 2.6. At first we show that f is a fuzzy continuous function respect to fuzzy metric M .

Assume that ϵ and t are arbitrary positive numbers and $\varphi : \mathbb{R}^+ \rightarrow (0, 1]$ is the map in Example 2.6. By investigating various conditions of x and y , we can prove that

$$\frac{\min\{f(x), f(y)\}}{\max\{f(x), f(y)\}} > \frac{1}{10} \frac{\min\{x, y\}}{\max\{x, y\}},$$

for all $x, y \in X$. For example if $x, y \in [0, \frac{1}{2}]$ and $x < y$ then $f(x) = \frac{3}{4}x + \frac{1}{8}$ and $f(y) = \frac{3}{4}y + \frac{1}{8}$. So $\frac{\min\{f(x), f(y)\}}{\max\{f(x), f(y)\}} = \frac{\frac{3}{4}x + \frac{1}{8}}{\frac{3}{4}y + \frac{1}{8}}$ and clearly $\frac{\frac{3}{4}x + \frac{1}{8}}{\frac{3}{4}y + \frac{1}{8}} > \frac{1}{10} \frac{x}{y}$.

This implies that $M(f(x), f(y), t) > M(x, y, \frac{t}{10})$. So this is clear that if $M(x, y, \frac{t}{10}) > 1 - \epsilon$ then $M(f(x), f(y), t) > 1 - \epsilon$. This prove that f is a fuzzy continuous

FIGURE 1. The graph of the function f from Example 4.3

function on $(X, M, *)$.

Let $\epsilon = \frac{1}{5}$. Since $f(0) = \frac{1}{8}$, $f(\frac{1}{2}) = \frac{1}{2}$ and $f(\frac{3}{4}) = \frac{7}{8}$, for any $\delta > 0$ we can find the points $a, b \in X$ such that $a < \frac{1}{2} < b$, $\frac{\min\{f(a), \frac{1}{2}\}}{\max\{f(a), \frac{1}{2}\}} > 1 - \delta$ and $\frac{\min\{f(\frac{1}{2}), b\}}{\max\{f(\frac{1}{2}), b\}} > 1 - \delta$.

Up to this point, there is a δ - F -pseudo orbit that contains $\frac{1}{4}$ and 1. Let $x \in X$, if $x \leq \frac{1}{2}$ then $M(f^i(x), \frac{1}{4}, t) < 1 - \frac{1}{5}$, for all $t > 1$ and for all $i \geq 0$. And, if $x < \frac{1}{2}$ then $M(f^i(x), 1, t) < 1 - \frac{1}{5}$, for all $t > 1$ and for all $i \geq 0$. This implies that $(X, M, *, f)$ does not have the F -shadowing property.

(d) Suppose $(X, M_1, *)$ is the fuzzy metric space in Example 2.7. Given $\epsilon > 0$, since f is continuous then there is $\delta > 0$ such that if $|y - x| < \delta$ then $|f(y) - f(x)| < \frac{\epsilon}{8}$. Let $x < y$ and $\frac{x}{y} > 1 - \delta$. This implies that $y - x < \delta$, so $f(y) - f(x) < \frac{\epsilon}{8}$ and so

$\frac{f(x)}{f(y)} > 1 - \frac{\epsilon}{8f(y)} > 1 - \epsilon$. Thus f is a fuzzy continuous function on $(X, M_1, *)$. By similar argument to the previous paragraph, one can prove that $(X, M_1, *, f)$ does not have the F -shadowing property.

It is a remarkable fact that, one can prove analogous results for any map g sufficiently close to f .

EXAMPLE 4.4. Suppose $\alpha < \frac{1}{128}$ is a positive number and $g : [0, 1] \rightarrow [0, 1]$ be a continuous monotone map such that $g(x) > x$ if and only if $x \in [0, \frac{1}{2}) \cup (\frac{1}{2}, 1)$, $g(1) = 1$, $g(\frac{1}{2}) = \frac{1}{2}$ and




FIGURE 2. The graph of the functions f and g from Example 4.3 that are near together.

$\sup\{|f(x) - g(x)| : x \in [0, 1]\} < \alpha$ (see Figure 2). At first we show that $(X, M_1, *, g)$ is a fuzzy continuous function. Since $(X, M_1, *, g)$ is a fuzzy continuous function, this is sufficient to prove that $M_1(g(x), g(y), t) > \frac{1}{2} M_1(f(x), f(y), t)$ for all $x, y \in X$ and $t \in [0, 1]$.

Let $x, y \in [0, 1]$ which $x < y$. It is clear that $f(x)f(y) > \alpha(f(x) + f(y))$. So $\frac{f(x)-\alpha}{f(y)+\alpha} > \frac{1}{2} \frac{f(x)}{f(y)}$. This implies that $\frac{g(x)}{g(y)} > \frac{f(x)-\alpha}{f(y)+\alpha} > \frac{1}{2} \frac{f(x)}{f(y)}$. Then

$$\frac{\min\{g(x), g(y)\}}{\max\{g(x), g(y)\}} = \frac{g(x)}{g(y)} > \frac{1}{2} \frac{f(x)}{f(y)} = \frac{1}{2} \frac{\min\{f(x), f(y)\}}{\max\{f(x), f(y)\}}.$$

So $M_1(g(x), g(y), t) > \frac{1}{2} M_1(f(x), f(y), t)$ for all $x, y \in X$ and $t \in [0, 1]$.

Similar argument to f in Example 4.3 shows that the map g does not have the shadowing property and $(X, M_1, *, g)$ does not have the F -shadowing property.

5. CONCLUSION

In this paper, we extend some notions and results known in discrete dynamical systems on fuzzy discrete dynamical systems and some related properties are investigated. In particular, the relation between various fuzzy shadowing property and fuzzy mixing is investigated. More precisely, the following implications are obtained:

(a) F -ergodic shadowing property implies F -chain transitivity.

- (b) F -ergodic shadowing property implies F -shadowing property.
 (c) F -shadowing and F -chain mixing properties imply F -topologically mixing property.

By definitions, most of the proofs were straight forward, but because of Example 4.3 we expect that a theorem similar to Theorem 2.11 in fuzzy discrete dynamical systems can be obtained for special fuzzy metric spaces. In particular, We are going to study and extend the other types of shadowing and their equivalence to fuzzy topology in another papers. Investigating the relation between chaos and stability with shadowing in fuzzy topology will become our future research topic. We finish this paper with the following important questions:

- 1) For a fuzzy discrete dynamical system $f : X \longrightarrow X$, are the F -ergodic shadowing and F -topological mixing properties equivalent?
- 2) Does the fuzzy discrete dynamical system $(X, M, *, f_\beta)$ have the F -shadowing and F -chain mixing properties, when f_β is the tent map and $(X, M_1, *)$ is the fuzzy metric space presented in Example 2.6 or Example 2.7?

ACKNOWLEDGMENTS

The author is thankful of referees for their valuable comments.

REFERENCES

1. R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms*, Springer Lecture Notes in Math, 1975.
2. J. S. Canovas, J. Kupka, Topological Entropy of Fuzzified Dynamical Systems, *Fuzzy Sets and Systems*, **165**, (2011), 37-49.
3. L. Chen, S. Li, Shadowing Property for Inverse Limit Spaces, *Trans. Amer. Math. Soc.*, **115** (2), (1992), 573-580.
4. E. M. Coven, I. Kan, J.A. Yorke, Pseudo-orbit Shadowing in the Family of Tent Maps. *Trans. Amer. Math. Soc.*, **308** (1), (1988), 227-241.
5. A. Fakhari, F. Helen Ghane, On Shadowing: Ordinary and Ergodic, *J. Math. Anal. Appl.*, **364** (1), (2010), 151-155.
6. M. Fatehi Nia, Parameterized IFS with the Asymptotic Average Shadowing Property, *Qual. Theory Dyn. Syst.*, **15**(2), (2016), 367-381.
7. M. Fatehi Nia, Iterated function systems with the average shadowing property, *Topology Proc.*, **48**, (2016), 261-275.
8. A. George, P. Veeramani, On some results of analysis for fuzzy metric spaces, *Fuzzy Sets and Systems*, **90** (3), (1997), 365-368.
9. A. George, P. Veeramani, On some results in fuzzy metric spaces. *Fuzzy Sets and Systems*, **64** (3), (1994), 395-399.
10. M. Grabiec, Fixed points in fuzzy metric spaces, *Fuzzy Sets and Systems*, **27**, (1988), 385-389.
11. V. Gregori, S. Morillas, A. Sapena, Examples of fuzzy metrics and applications, *Fuzzy Sets and Systems*, **170**, (2011), 95-111.
12. V. Gregori, S. Romaguera, Some properties of fuzzy metric spaces, *Fuzzy Sets and Systems*, **115**(3), (2000), 485-489.

13. R. Gu, *Shadowing Techniques and Chaotic Phenomena*, In Advances in Discrete Dynamics, Nova Science Publishers, New York, 65-90, 2012.
14. R. Gu, The asymptotic average shadowing property and transitivity, *Nonlinear Analysis* **67**, (2007), 1680-1689.
15. I. Kramosil, J. Michalek, Fuzzy metrics and statistical metric spaces, *Kybernetika (Prague)*, **11**(5), (1975), 336-344.
16. J. Kupka, On devaney chaotic induced dynamical systems, *Fuzzy Sets and Systems*, **177**, (2011), 34-44.
17. J. Kupka, On fuzzifications of discrete dynamical systems, *Inform Sciences*, **181**, (2011), 2858-2872.
18. D. Mihet, A Banach contraction theorem in fuzzy metric spaces. *Fuzzy Sets and Systems*, **144**(3), (2004), 431-439.
19. K. Palmer, *Shadowing in dynamical systems: theory and applications*, Kluwer Acad. Publ., Dordrecht, 2000.
20. D. Richeson, J. Wiseman, Chain Recurrence Rates and Topological Entropy, *Topology Appl.* **156**, (2008), 251-261.
21. J. Rodriguez-Lopez, S. Romaguera, The Hausdorff fuzzy metric on compact sets, *Fuzzy Sets and Systems*, **147**, (2004), 273-283.
22. H. Roman-Flores, Y. Chalco-Cano, Some Chaotic Properties of Zadeh's Extensions, *Chaos Solitons and Fractals*, **35**(3), (2008), 452-459.
23. K. Sakai, Various Shadowing Properties for Positively Expansive Maps, *Topology. Appl.* **131** (10), (2003), 15-31.
24. S. Sedaghat, Y. Ordokhani, Stability and numerical solution of time variant linear systems with delay in both the state and control, *Iran. J. Math. Sci. Inform.*, **7**, (2012), 43-57.
25. P. Walters, On the Pseudo-orbit Tracing Property and its Relationship to Stability, *Lecture Notes in Math, Springer-Verlag*, **668**, (1978), 231-244.
26. R. S. Yang, The Pseudo-orbit Tracing Property and Chaos, *Acta Math. Sinica.*, **39**, (1996), 382-386.
27. R. S. Yang, Pseudo-orbit tracing property and completely positive entropy, *Acta Math. Sinica.*, **42**, (1999), 99-104.
28. M. Yarahmadi, S.M. Karbassi, Design of robust controller by neuro-fuzzy system in a prescribed region via state feedback, *Iran. J. Math. Sci. Inform.*, **4**, (2009), 1-16.