Iranian Journal of Mathematical Sciences and Informatics

Vol. 14, No. 2 (2019), pp 153-157

DOI: 10.7508/ijmsi.2019.02.012

Copresented Dimension of Modules

M. Amini*, F. Hassani

Department of Mathematics, Payame Noor University, Tehran, Iran.

E-mail: mostafa.amini@pnu.ac.ir E-mail: Hassani@pnu.ac.ir

ABSTRACT. In this paper, a new homological dimension of modules, copresented dimension, is defined. We study some basic properties of this homological dimension. Some ring extensions are considered, too. For instance, we prove that if $S \geq R$ is a finite normalizing extension and S_R is a projective module, then for each right S-module M_S , the copresented dimension of M_S does not exceed the copresented dimension of $Hom_R(S, M)$.

Keywords: Coherent ring, Copresented dimension, Projective module.

2000 Mathematics Subject Classification: 13D07, 16D40, 18G25

1. Introduction

Throughout this paper, R is an associative ring with identity and all modules are unitary. First we recall some known notions and facts needed in the sequel. Let R be a ring, n a non-negative integer and M an R-module. Then

- (1) M is said to be *finitely cogenerated* [1] if for every family $\{V_k\}_J$ of submodules of M with $\bigcap_J V_k = 0$, there is a finite subset $I \subset J$ with $\bigcap_I V_k = 0$.
- (2) M is said to be n-copresented [14] if there is an exact sequence of R-modules $0 \to M \to E^0 \to E^1 \to \cdots \to E^n$, where each E^i is a finitely cogenerated injective module.

Received 06 March 2017; Accepted 03 March 2018 ©2019 Academic Center for Education, Culture and Research TMU

^{*}Corresponding Author

Downloaded from ijmsi.com on 2025-10-16

- (3) R is called right *co-coherent* [17] if every finitely cogenerated factor module of a finitely cogenerated injective R-module is finitely copresented.
- (4) R is called n-cocoherent [14] in case every n-copresented R-module is (n+1)-copresented. It is easy to see that R is cocoherent if and only if it is 1-cocoherent. Recall that a ring R is called right conoethrian [4] if every factor module of a finitely cogenerated R-module is finitely cogenerated. By [4, Proposition 17], a ring R is co-noethrian if and only if it is 0-cocoherent.
- (5) M is said to be n-presented [5] if there is an exact sequence of R-modules $F_n \to F_{n-1} \to \cdots \to F_1 \to F_0 \to M \to 0$, where each F_i is a finitely generated free module.
- (6) R is called *coherent* [18] in case every 0-presented R-module is 1-presented.
- (7) A ring extension $R \subseteq R'$ with characteristic p > 0 is called a *purely inseparable extension* [10] if for every element $r' \in R'$, there exists a non-negative integer n such that $r'^{p^n} \in R$.
- (8) For any commutative ring R of prime characteristic p > 0, assume that $F_R : R \to R^{(e)}$ is the e-th iterated Frobenius map in which $R^{(e)} \cong R$. Then, the *perfect closure* [9] of R, denoted by R^{∞} , is defined as the limit of the following direct system:

$$R \xrightarrow{F_R} R \xrightarrow{F_R} R \xrightarrow{F_R} \cdots$$

- (9) M is called (n,d)-injective [18] if $\operatorname{Ext}_R^{d+1}(N,M)=0$ for any n-presented right R-module N. It is clear that M is (0,0)-injective if and only if M is injective.
- (10) Assume that $S \geq R$ is a unitary ring extension. Then, the ring S is called right R-projective [6] in case, for any right S-module M_S with an S-module N_S , $N_R \mid M_R$ implies $N_S \mid M_S$, where $N \mid M$ means that N is a direct summand of M.
- (11) The ring extension $S \geq R$ is called a *finite normalizing extension* [8] in case there is a finite subset $\{s_1, \dots, s_n\} \subseteq S$ such that $S = \sum_{i=1}^{i=n} s_i R$ and $s_i R = Rs_i$ for $i = 1, \dots, n$.
- (12) A finite normalizing extension $S \geq R$ is called an almost excellent extension [12] in case RS is flat, SR is projective, and the ring S is right R-projective.

In this paper, we introduce the dual concepts of presented dimensions of R-modules. We also, introduce the copresented dimension of any R-module M:

 $\operatorname{FEd}(M) = \inf\{m \mid \text{ there exists an injective resolution } 0 \to M \to E^0 \to \cdots \to E^m \to \cdots \to E^{m+i} \to \cdots$, such that E^{m+i} are finitely cogenerated for

 $i=0,1,2,\cdots$. If $K=\ker(E^m\to E^{m+1})$, then K has an infinite finite copresentation. It is clear that any copresented dimension is finitely copresented dimension (see [16]). Also, the copresented dimension of ring R is defined to be:

$$FED(R) = \sup\{FEd(M) \mid M \text{ is a finitely cogenerated module}\}.$$

Then, some basic properties of the copresented dimensions of modules are studied. For example, it is shown that if $\operatorname{FEd}(M) < \infty$, then $\operatorname{id}(M) \le n$ if and only if $\operatorname{Ext}_R^{n+1}(N,M) = 0$ for every strongly copresented R-module N. Also, it is proved that $\operatorname{FED}(R \oplus S) = \sup\{\operatorname{FED}(R), \operatorname{FED}(S)\}$, for any two rings R and S. Also, some characterizations of the copresented dimensions of modules on Ring Extensions are determined. For instance, let $S \ge R$ be a finite normalizing extension with S_R projective as an R-module, then for any right R-module M_R , we have $\operatorname{FEd}(\operatorname{Hom}_R(S,M))_S \le \operatorname{FEd}(M_R)$. Finally, we give a sufficient condition under which $\operatorname{FED}(S) \le \operatorname{FED}(R)$ and or $\operatorname{FED}(R) < \operatorname{FED}(S) + \max\{k,d\}$, where $k = \operatorname{id}(S_R)$ and $d = \sup\{\operatorname{FEd}(M_R) \mid M \in \operatorname{Mod} - S$ and $\operatorname{FEd}(M_S) = 0\}$.

2. Main Results

We start this section with the following definition which is the dual of the presented dimension of a module.

Definition 2.1. For any R-module M, we define the copresented dimension of M to be $\operatorname{FEd}(M) = \inf\{m \mid \text{there exists an injective resolution } 0 \to M \to E^0 \to \cdots \to E^m \to \cdots \to E^{m+i} \to \cdots$, so that E^{m+i} are finitely cogenerated for $i=0,1,2,\cdots\}$. In particular, a module M is called strongly copresented module if $\operatorname{FEd}(M) = 0$.

Proposition 2.2. For any R-module M, $FEd(M) \leq id(M) + 1$.

Proof. It is a direct consequence of Definition 2.1.

EXAMPLE 2.3. Let $R = \mathbb{Z}$. Since $\mathrm{id}(\mathbb{Z}_{p^{\infty}}) = 0$, we have $\mathrm{FEd}(\mathbb{Z}_{p^{\infty}}) \leq 1$. On the other hand, $\mathbb{Z}_{p^{\infty}}$ is finitely cogenerated by [1, p.124]. So by Definition 2.1, $\mathrm{FEd}(\mathbb{Z}_{p^{\infty}}) = 0$.

Now, we study the behavior of the copresented dimension on the exact sequences. Before this we need the following lemma.

Lemma 2.4. Let $0 \to A \xrightarrow{f'} B \xrightarrow{f} C \to 0$ be a short exact sequence of R-modules. Then:

(1) If $0 \to A \to A^0 \to A^1 \to \cdots$ and $0 \to C \to C^0 \to C^1 \to \cdots$ are injective resolutions of A and C, respectively. Then the exact sequence $0 \longrightarrow B \longrightarrow A^0 \oplus C^0 \longrightarrow A^1 \oplus C^1 \longrightarrow \cdots$

is an injective resolution of B.

(2) If $0 \to B \to B^0 \to B^1 \to \cdots$ and $0 \to C \to C^0 \to C^1 \to \cdots$ are injective resolutions of B and C, respectively. Then the exact sequence

$$0 \longrightarrow A \longrightarrow B^0 \longrightarrow D^0 \longrightarrow D^1 \longrightarrow \cdots$$

is an injective resolution of A, where $D^i = C^i \oplus B^{i+1}$ for any $i \geq 0$.

(3) If $0 \to B \to B^0 \to B^1 \to \cdots$ and $0 \to A \to A^0 \to A^1 \to \cdots$ are injective resolutions of B and A, respectively. Then the exact sequence

$$0 \longrightarrow C \longrightarrow F^0 \longrightarrow E^0 \longrightarrow E^1 \longrightarrow \cdots$$

is an injective resolution of C, where $F^0 = B^0 \oplus A^1$ and $E^i = A^0 \oplus B^{i+1} \oplus A^{i+2}$ for any $i \geq 0$.

Proof. (1) The proof is similar to that of [3, Theorem 2.4].

(2) Let $0 \to B \to B^0 \to B^1 \to \cdots$ be an injective resolution of B. Then, the exact sequences

 $0 \to K \to B^1 \to B^2 \to \cdots$ and $0 \to B \to B^0 \to K \to 0$ exist, where $K = \frac{B^0}{B}$. Now, we consider the following commutative diagram:

By (1), there is an exact sequence

$$0 \longrightarrow D \longrightarrow D^0 \longrightarrow D^1 \longrightarrow D^2 \longrightarrow \cdots$$

of injective R-modules D^i such that $D^i = C^i \oplus B^{i+1}$ for any $i \geq 0$.

Combining this sequence with the exact sequence $0 \to A \to B^0 \to D \to 0$, we get the exact sequence

$$0 \longrightarrow A \longrightarrow B^0 \longrightarrow D^0 \longrightarrow D^1 \longrightarrow \cdots$$

where B^0 and D^i are injective for any $i \geq 0$.

(3) Let $0 \to A \to A^0 \to A^1 \to \cdots$ be an injective resolution of A. Then, the exact sequences

 $0 \to K \to A^1 \to A^2 \to \cdots$ and $0 \to A \to A^0 \to K \to 0$ exist, where $K = \frac{A^0}{A}$. Now, we consider the following commutative diagram:

By (1), there is an exact sequence

$$0 \longrightarrow F \longrightarrow F^0 \longrightarrow F^1 \longrightarrow F^2 \longrightarrow \cdots$$

of injective R-modules F^i such that $F^i = B^i \oplus A^{i+1}$ for any $i \ge 0$.

It is clear that $F = A^0 \oplus C$. So, the exact sequence $0 \to C \to F \to A^0 \to 0$ exists. Let $K = \frac{F^0}{F}$, then we obtain the following commutative diagram:

Therefore by (1), the sequence

$$0 \longrightarrow E \longrightarrow E^0 \longrightarrow E^1 \longrightarrow E^2 \longrightarrow \cdots$$

is an injective resolution of E, where $E^i = A^0 \oplus F^{i+1} = A^0 \oplus B^{i+1} \oplus A^{i+2}$ for any $i \geq 0$. Combining this sequence with the exact sequence $0 \to C \to F^0 \to E \to 0$, we get the exact sequence

$$0 \longrightarrow C \longrightarrow F^0 \longrightarrow E^0 \longrightarrow E^1 \longrightarrow \cdots$$

where F^0 and E^i are injective for any i > 0.

Theorem 2.5. Let $0 \to A \xrightarrow{g} B \xrightarrow{f} C \to 0$ be an exact sequence of R-modules. Then $FEd(B) \le \max\{FEd(A), FEd(C)\}, FEd(C) \le \max\{FEd(B), FEd(A) + 1\}, FEd(A) \le \max\{FEd(B), FEd(C) - 1\}.$

Proof. Assume that $\mathbf{E}^{'}$ is an injective resolution of A and $\mathbf{E}^{''}$ is an injective resolution of C. Thus by Lemma 2.5(1), there exists an injective resolution \mathbf{E} of B such that

$$0 \to \mathbf{E}'^{\mathbf{A}} \to \mathbf{E}^{\mathbf{B}} = \mathbf{E}'^{\mathbf{A}} \oplus \mathbf{E}''^{\mathbf{C}} \to \mathbf{E}''^{\mathbf{C}} \to 0$$

is an exact sequence of complexes. Hence for every $m \ge \max\{\operatorname{FEd}(A),\operatorname{FEd}(C)\}$, E^m is finitely cogenerated. So, we deduce that $\operatorname{FEd}(B) \le \max\{\operatorname{FEd}(A),\operatorname{FEd}(C)\}$.

Assume that \mathbf{E}'' is an injective resolution of C and \mathbf{E} is an injective resolution of B. Thus by Lemma 2.5(2), the exact sequence

$$0 \longrightarrow A \longrightarrow E^0 \longrightarrow D^0 \longrightarrow D^1 \longrightarrow \cdots \longrightarrow D^d \longrightarrow \cdots$$

is an injective resolution of A. So for every $d \ge \max\{\operatorname{FEd}(B), \operatorname{FEd}(C) - 1\}$, D^d is finitely cogenerated. Thus, we have that $\operatorname{FEd}(A) \le \max\{\operatorname{FEd}(B), \operatorname{FEd}(C) - 1\}$. Also, it is prove that $\operatorname{FEd}(C) \le \max\{\operatorname{FEd}(B), \operatorname{FEd}(A) + 1\}$.

The proof of the following Corollary is similar to the proof of [19, Corollary 2.7].

Corollary 2.6. If $FEd(M_1)$, $FEd(M_2)$, \cdots $FEd(M_d)$ are finite, then:

$$FEd(\oplus M_i) = \max\{FEd(M_i) \mid i = 1, \dots, d\}.$$

Proof. For the case m=2, the exact sequences

$$0 \to M_1 \to M_1 \oplus M_2 \to M_2 \to 0$$

and

$$0 \to M_2 \to M_2 \oplus M_1 \to M_1 \to 0$$

exist. Thus by Theorem 2.5, we deduce that

$$\operatorname{FEd}(M_2) \leq \max\{\operatorname{FEd}(M_1 \oplus M_2), \operatorname{FEd}(M_1) - 1\},\$$

$$FEd(M_1) \le \max\{FEd(M_1 \oplus M_2), FEd(M_2) - 1\}$$

and

$$FEd(M_1 \oplus M_2) \le \max\{FEd(M_1), FEd(M_2)\}.$$

Assume that $FEd(M_1) < FEd(M_2)$. Then $FEd(M_1) \le FEd(M_2) - 1$, and we have:

$$FEd(M_2) \le \max\{FEd(M_1 \oplus M_2), FEd(M_2) - 2\} = FEd(M_1 \oplus M_2).$$

Also, similarly $\text{FEd}(M_1) \leq \text{FEd}(M_1 \oplus M_2)$. So, we conclude that $\text{FEd}(M_1 \oplus M_2) = \max\{\text{FEd}(M_1), \text{FEd}(M_2)\}$.

Proposition 2.7. Let n be a non-negative integer. Then the following statements are equivalent:

- (1) $id(M) \le n$ for every strongly corresented R-module M;
- (2) $\operatorname{Ext}_R^{n+1}(N,M) = 0$ for every strongly copresented R-module N.

Proof. $(1) \Rightarrow (2)$ This is obvious.

 $(2) \Rightarrow (1)$ We use the induction on n. Let n = 0. Since $\operatorname{Ext}_R^1(N, M) = 0$ for any strongly copresented R-module N, by using the exact sequence $0 \to M \to E^0 \to L^0 \to 0$ where E^0 is finitely cogenerated and L^0 is strongly copresented, we deduce that $\operatorname{Ext}_R^1(L^0, M) = 0$. Therefore by [7, Theorem 7.31], the exact sequence ebove is split. So, M is injective and hence $\operatorname{id}(M) < 0$. Assume that

Downloaded from ijmsi.com on 2025-10-16

n>0. By [7, Corollary 6.42], we have that $\operatorname{Ext}_R^{n+1}(N,M)\cong\operatorname{Ext}_R^n(N,L^0)=$ 0. Thus by induction hypothesis, $id(L^0) \leq n-1$. Therefore from the exact sequence ebove, we deduce that $id(M) \leq n$.

Proposition 2.8. Let $FEd(M) \leq 1$. Then the following statements are equivalent:

- (2) Ext_Rⁿ⁺¹(N, M) = 0 for every strongly copresented R-module N.

Proof. Since $FEd(M) \leq 1$, the exact sequence $0 \to M \to E^0 \to L^0 \to 0$ exists, where E^0 is injective and L^0 is strongly copresented. Thus, $\operatorname{Ext}_R^{n+1}(N,M)=0$ for any strongly copresented R-module N if and only if $\operatorname{Ext}_R^n(N, L^0) = 0$ if and only if $id(L^0) \le n-1$ (by Proposition 2.7) if and only if $id(M) \le n$.

Theorem 2.9. Let $FEd(M) < \infty$. Then the following statements are equivalent:

- $\begin{array}{ll} (1) \ \operatorname{id}(M) \leq n; \\ (2) \ \operatorname{Ext}_R^{n+1}(N,M) = 0 \ for \ every \ strongly \ copresented \ R\text{-}module \ N. \end{array}$

Proof. $(1) \Rightarrow (2)$ It is clear.

 $(2) \Rightarrow (1)$ If FEd(M) = m, then the exact sequence

$$0 \to M \to E^0 \to E^1 \to \cdots \to E^{m-1} \overset{d^{m-1}}{\to} E^m \overset{d^m}{\to} \cdots \to E^{m+j} \to \cdots$$

exists, where E^i is finitely cogenerated for any $i \geq m$. By Proposition 2.2, $n+1 \geq m$. Let $\operatorname{Ext}_R^{n+1}(N,M) = 0$ for every strongly corresented R-module N. Thus by [7, Corollary 6.42], we have

$$\operatorname{Ext}_R^{n+1}(N,M) \cong \operatorname{Ext}_R^{n-m+1}(N,\operatorname{coker} d^{m-1}) = 0.$$

Since $\operatorname{coker} d^{m-1}$ is strongly copresented, Proposition 2.8 impleis that

$$id(coker d^{m-1}) \le n - m$$

and so, we deduce that $id(M) \leq n$.

Corollary 2.10. Let $D(R) < \infty$. Then:

$$D(R) = \sup\{pd(N) \mid N \text{ is strongly copresented}\}.$$

Proof. Assume that D(R) < m. Thus, pd(N') < m for any R-module N'. So, for any strongly corresented R-module N, $pd(N) \leq m$. Conversely, let $\operatorname{pd}(N) \leq m$ for every strongly corresented R-module N. Thus $\operatorname{Ext}_{R}^{m+1}(N,M) =$ 0 for every strongly presented R-module M. Since $D(R) < \infty$, $FEd(M) < \infty$ by Proposition 2.2. Therefore by Theorem 2.9, $id(M) \leq m$ and hence by [19, corollary 3.7], $D(R) \leq m$.

Definition 2.11. For any ring R, we define the corresented dimension of Rto be $FED(R) = \sup\{FEd(M) \mid M \text{ is a finitely cogenerated module}\}.$

EXAMPLE 2.12. Let $R = k[x^3, x^3y, xy^3, y^3]$, where k is a field with characteristic p = 3. By Definition 2.11 and Proposition 2.2, FED $(R^{\infty}) \leq D(R^{\infty}) + 1$, where R^{∞} is perfect closure of R. On the other hand, k[x, y] is purely inseparable over R. Also, by [9, Proposition 3.3], $(k[x, y])^{\infty}$ is coherent. Therefore by [10, Remark 1.4], R^{∞} is coherent. Since R is reduced, [2, Proposition 5.5] implies that FED $(R^{\infty}) \leq \dim(R) + 1$ and so, FED $(R^{\infty}) \leq 3$.

Proposition 2.13. The following statements are equivalent:

- (1) FED(R) = 0;
- (2) Every finitely cogenerated module has an infinite finite copresented;
- (3) Every finitely cogenerated module is finitely copresented;
- (4) R is co-noetherian.

Proof. The implication $(1) \Longrightarrow (2) \Longrightarrow (3)$ follow immediately from Definiton 2.11.

$$(3) \Longrightarrow (4) \Longrightarrow (1)$$
 are trivial.

Corollary 2.14. If $FED(R) \leq 0$, then R is n-cocoherent.

Proof. Since every n-copresented module M is finitely cogenerated, Proposition 2.13 implies that M is (n + 1)-copresented.

Next, we study the copresented dimension of the direct sum of rings. But before this we need the following lemma.

Lemma 2.15. Let $f: R \to S$ be a ring epimorphism. If M_S is a right S-module (hence a right R-module) and N_R is a right R-module, then the following statements hold:

- (1) $M \otimes_R S \cong M_S$.
- (2) If f is flat and N_R is a finitely cogenerated right R-module, then $N \otimes_R S$ is a finitely cogenerated right S-module.
- (3) If f is flat, then M_S is a finitely cogenerated right S-module if and only if M_R is a finitely cogenerated right R-module.
- (4) If f is projective, then M_S is an injective right S-module if and only if M_R is an injective right R-module.

Proof. (1) This is clear.

- (2) For any family of submodules $\{N_i \otimes_R 1_S | i \in I\}$ in $N \otimes_R S$, if $\bigcap (N_i \otimes_R 1_S) = 0$, then we need to show that $\bigcap_{i \in F} (N_i \otimes_R 1_S) = 0$ for some finite subset F of I. Since f is flat, we have that $\bigcap_{i \in I} N_i \otimes_R 1_S = 0$. So, $\bigcap_{i \in I} N_i = 0$ and hence by hypotises $\bigcap_{i \in F} N_i = 0$ for some finite subset F of I. Therefore, $\bigcap_{i \in F} (N_i \otimes_R 1_S) = \bigcap_{i \in F} N_i \otimes_R 1_S = 0$.
- (3) (\Rightarrow): Let $\psi: M \to \prod_{i \in I} R$ is a monomorphism, then we claim that $\pi: M \to \prod_{i \in F} R$ is a monomorphism for some finite subset F of I. We have the following commutative diagram:

$$\begin{array}{ccc}
M & \xrightarrow{\psi} & \prod_{i \in I} R \\
\downarrow \cong & & \downarrow g \\
M & \xrightarrow{h} & \prod_{i \in I} S,
\end{array}$$

where since g is epimorphism and ψ is monomorphism, h is monomorphism. So by hypothesis, $\alpha: M \to \prod_{i \in F} S$ is a monomorphism for some finite subset F of I. Therefore the following commutative diagram:

$$\begin{array}{ccc} M & \stackrel{\gamma}{\longrightarrow} & \prod_{i \in F} R \\ \downarrow \cong & & \downarrow \beta \\ M & \stackrel{\alpha}{\longrightarrow} & \prod_{i \in F} S, \end{array}$$

where β is epimorphism and α is monomorphism, implies that γ is monomorphism.

 (\Leftarrow) : This follows from (1) and (2)

(4) By [5, Lemma 3.3], M_S is an (n, d)-injective right S-module if and only if M_R is an (n, d)-injective right R-module. If n = 0, d = 0, Then (4) is hold. \square

Theorem 2.16. Assume that R and S are two rings. Then:

$$FED(R \oplus S) = \sup\{FED(R), FED(S)\}.$$

Proof. We first show that $\text{FED}(R \oplus S) \leq \sup\{\text{FED}(R), \text{FED}(S)\}$. Consider FED(R) = n, FED(S) = m and $n \geq m$. Also, let M be a finitely cogenerated right $(R \oplus S)$ -module. Then M has a unique decomposition $M = A \oplus B$, where A, B are right modules of rings R and S, respectively. By [15, Lemma 1.1], A and B are finitely cogenerated right $(R \oplus S)$ -module. So by Lemma 2.15, A is finitely cogenerated right R-module and B is finitely cogenerated right S-module. Therefore $\text{FEd}(A) \leq n$ and $\text{FEd}(B) \leq m$, and hence there is an exact sequences

$$0 \to A \to E_a^0 \to E_a^1 \to \cdots \to E_a^{n-1} \to E_a^n \to \cdots,$$

$$0 \to B \to E_b^0 \to E_b^1 \to \cdots \to E_b^{m-1} \to E_b^m \to \cdots$$

of injective right R-modules E_a^i and injective right S-modules E_b^i such that E_a^i, E_b^i are finitely cogenerated for any $i \geq n$ and $i \geq m$, respectively. So, we deduce that the exact sequence

$$0 \to A \oplus B \to E_a^0 \oplus E_b^0 \to E_a^1 \oplus E_b^1 \to \cdots \to E_a^{n-1} \oplus E_b^{m-1} \to E_a^n \oplus E_b^m \to \cdots$$

exists, where by Lemma 2.15, every $E_a^i \oplus E_b^i$ is injective right $(R \oplus S)$ -module and also, every $E_a^i \oplus E_b^i$ is finitely cogenerated for any $i \geq n$. Therefore, we have $\text{FED}(R \oplus S) \leq \sup\{\text{FED}(R), \text{FED}(S)\}$.

Conversely, Assume that $\text{FED}(R \oplus S) = d$. If M is a finitely cogenerated right R-module. Then by Lemma 2.15, M is a finitely cogenerated right $(R \oplus S)$ -module and hence $\text{FED}(M_{(R \oplus S)}) \leq d$. Thus, the exact sequence $0 \to M \to E^0 \to E^1 \to \cdots \to E^{d-1} \to E^d \to \cdots$ of injective right $(R \oplus S)$ -modules

 E^i exists, where every E^i is finitely cogenerated for any $i \geq d$. Let $E^i = C^i \oplus D^i$, where C^i is a R-module and D^i is a S-module. On the other hand, M is a right R-module, so we have the exact sequence $0 \to M \to C^0 \to C^1 \to \cdots \to C^{d-1} \to C^d \to \cdots$ of R-modules. But, every C^i is injective right $(R \oplus S)$ -module and also every C^i is finitely cogenerated right $(R \oplus S)$ -module for $i \geq d$. So by [15, Lemma 1.1] and Lemma 2.15, C^i is an injective right R-module and it is finitely cogenerated R-module for $i \geq d$. Therefore $FEd(M) \leq d$ and hence $FED(R) \leq d$. Similarly, $FED(S) \leq d$ and implies that $\sup\{FED(R), FED(S)\} \leq FED(R \oplus S)$.

Proposition 2.17. Let $S \geq R$ be a finite normalizing extension with S_R projective as an R-module. Then for any right R-module M_R , $\text{FEd}(\text{Hom}_R(S, M))_S \leq \text{FEd}(M_R)$.

Proof. Asume that $FEd(M_R) = n$. Then there axists an exact sequence of injective R-modules

$$0 \to M \to E^0 \to E^1 \to \cdots \to E^{n-1} \to E^n \to \cdots$$

where each E^i is finitely cogenerated for any $i \geq n$. Since S is projective, there is an exact sequence

$$0 \to \operatorname{Hom}_R(S, M) \to \operatorname{Hom}_R(S, E^0) \to \cdots \to \operatorname{Hom}_R(S, E^n) \to \cdots$$

of injective S-modules $\operatorname{Hom}_R(S, E^i)$, where by [13, Propositon 8.3], $\operatorname{Hom}_R(S, E^i)$ is finitely cogenerated for any $i \geq n$. Thus $\operatorname{FEd}(\operatorname{Hom}_R(S, M))_S \leq n$ and hence, we have $\operatorname{FEd}(\operatorname{Hom}_R(S, M))_S \leq \operatorname{FEd}(M_R)$.

Proposition 2.18. Let $S \geq R$ be a finite normalizing extension, S_R be Projective, and S be R-projective. Then for each right S-module M_S , $FEd(M_S) \leq FEd(Hom_R(S, M))$.

Proof. By [12, Lemma 1.1], M_S is isomorphic to a direct summand of $\operatorname{Hom}_R(S, M)$. So, from Corollary 2.6, we deduce that $\operatorname{FEd}(M_S) \leq \operatorname{FEd}(\operatorname{Hom}_R(S, M))$.

Proposition 2.19. Let $S \geq R$ be an almost excellent extension. Then for each right S-module M_S , $\text{FEd}(M_R) \leq \text{FEd}(M_S)$.

Proof. Asume that $FEd(M_S) = n$. So, there axists an exact sequence of injective S-modules

$$0 \to M \to E^0 \to E^1 \to \cdots \to E^{n-1} \to E^n \to \cdots$$

where each E^i is finitely cogenerated for any $i \geq n$. Thus by [18, Proposition 5.1], every E^i is an injective R-module and also, it is a finitely cogenerated R-module for $i \geq n$ by [14, Theorem 5]. Therefore, it follows that $FEd(M_R) \leq FEd(M_S)$.

Corollary 2.20. Let $S \geq R$ be an almost excellent extension. Then for each right S-module M_S , $FEd(M_R) = FEd(M_S) = FEd(Hom_R(S, M))$.

[Downloaded from ijmsi.com on 2025-10-16]

Theorem 2.21. Asume that $S \geq R$ is a finite normalizing extension and S_R is Projective. Then:

- (1) If S is R-projective and $FED(S) < \infty$, then $FED(S) \le FED(R)$.
- (2) If $FED(R) < \infty$, then $FED(R) < FED(S) + \max\{k, d\}$, where $k = id(S_R)$ and $d = \sup\{FEd(M_R) \mid M \in \text{Mod} S \text{ and } FEd(M_S) = 0\}$.

Proof. (1) Asume that FED(S) = n and $FEd(M_S) = n$ for a finitely cogenerated S-module M. Since S_R is projective, by hypothesis and [12, Lemma 1.1], M_S is isomorphic to a direct summand of $Hom_R(S, M)$ and hence we have:

$$0 \to K \to \operatorname{Hom}_R(S, M)) \to M_S \to 0.$$

By [14, Lemma 4], $\operatorname{Hom}_R(S, M)$) is finitely cogenerated S-module, since M_R is a finitely cogenerated R-module. So, $\operatorname{FEd}(\operatorname{Hom}_R(S, M)_S) \leq n$. On the other hand, by Theorem 2.5,

$$FEd(K) \le \max\{n, n-1\},\$$

 $n = \text{FEd}(M_S) \le \max\{\text{FEd}(\text{Hom}_R(S, M)_S), \text{FEd}(K_S) - 1\} \le \text{FED}(S) = n.$

Therefore $FEd(Hom_R(S, M)_S) = n$. Thus, Proposition 2.17 implies that

$$\operatorname{FEd}(\operatorname{Hom}_R(S, M)_S) \le \operatorname{FEd}(M_R)$$

and hence $FED(S) \leq FED(R)$.

(2) Assume that FED(R) = n and $FEd(M_R) = n$ for a finitely cogenerated R-module M. Since S_R is projective, by [12, Lemma 1.1], M_R is isomorphic to a direct summand of $Hom_R(S, M)$ which induces the following short exact sequence of R-modules:

$$0 \to K \to \operatorname{Hom}_R(S_R, M)) \to M_R \to 0.$$

It is clear that $\operatorname{Hom}_R(S_R,M)$) is a finitely cogenerated R-module. Thus Theorem 2.5 implies that

 $n = \text{FEd}(M_R) \le \max\{\text{FEd}(\text{Hom}_R(S_R, M)), \text{FEd}(K_R) - 1\} \le \text{FED}(R) = n,$ and hence $\text{FEd}(\text{Hom}_R(S_R, M)) = n.$

If $FEd(Hom_R(S, M))_S = m \le FED(S)$, then there is an injective resolution

$$0 \longrightarrow \operatorname{Hom}_{R}(S, M) \xrightarrow{f_{0}} E^{0} \xrightarrow{f_{1}} E^{1} \longrightarrow \cdots \longrightarrow E^{m-1} \xrightarrow{f_{m}} E^{m} \xrightarrow{f_{m+1}} \cdots$$

of $\operatorname{Hom}_R(S,M)$, where every E^i is a finitely cogenerated S-module for any $i \geq m$. Let $D^i = \operatorname{coker}(f_i)$ for every $i \geq 0$. Thus, the following short exact sequences

$$0 \longrightarrow \operatorname{Hom}_R(S,M) \longrightarrow E^0 \to D^0 \longrightarrow 0,$$

. . .

$$0 \longrightarrow D^{m-2} \longrightarrow E^{m-1} \longrightarrow D^{m-1} \longrightarrow 0,$$
$$0 \longrightarrow D^{m-1} \longrightarrow E^m \longrightarrow D^m \longrightarrow 0$$

exists, where $FEd(D^{m-1}) = 0$. But by hypothesis and Proposition 2.2, we have:

$$FEd(D^i)_R \le id(D^i)_R + 1 \le id(S_R) + 1 = k + 1$$
, $FEd(D^{m-1})_R \le d$.

Therefore by Theorem 2.5, we deduce that:

 $\operatorname{FEd}(D^{m-2})_R \leq \max\{\operatorname{FEd}(E^{m-1})_R, \operatorname{FEd}(D^{m-1})_R + 1\} < \max\{k+1, d+1\} = 1 + \max\{k, d\},$

$${\rm FEd}(D^{m-3})_R \leq \max\{{\rm FEd}(E^{m-2})_R, {\rm FEd}(D^{m-2})_R + 1\} < 2 + \max\{k,d\},$$

:

$$\begin{split} & \operatorname{FEd}(D^0)_R \leq \max\{\operatorname{FEd}(E^1)_R,\operatorname{FEd}(D^1)_R+1\} < m-1+\max\{k,d\}, \\ & n = \operatorname{FEd}(\operatorname{Hom}_R(S,M))_R \leq \max\{\operatorname{FEd}(E^0)_R,\operatorname{FEd}(D^0)_R+1\} < m+\max\{k,d\}. \\ & \operatorname{Thus} \ \operatorname{FED}(R) < m+\max\{k,d\} \leq \operatorname{FED}(S)+\max\{k,d\} \ \text{and so, the proof is complete.} \end{split}$$

Corollary 2.22. Let $S \ge R$ be an almost excellent extension. Then $FED(R) < FED(S) + id(S)_R$.

Proof. By Proposition 2.19 and Theorem 2.21, this is clear.
$$\Box$$

ACKNOWLEDGMENTS

The authors wish to thank referees for their valuable comments.

References

- F. W. Anderson, K. R. Fuller, Rings and Categories of Modules, New York, Spring-Verlag, 1992.
- M. Asgharzadeh, K. Shimomoto, Almost Cohen-Macaulay and Almost Regular Algebras via Almost Flat Extensions, J. Commut. Algebra, 4(4), (2012), 445–478.
- D. Bennis, H. Bouzraa, Abdul-Qawe Kaed, On n-Copresented Modules and n-Cocoherent Ring, Int. Electron. J. Algebra, 12(2012), 162-174.
- V. A. Hiremath, Cofinitely Generated and Cofinitely Related Modules, Acta Math. Hungar, 39 (1-3), (1982), 1-9.
- B. Ouyang, L. Duan, W. Li, Relative Projective Dimensions, Bull. Malays. Math. Sci. Soc., (2)37(3), (2014), 865-879.
- D. S. Passman, The Algebraic Structure of Group Rings, John Wiley and Sons, New York, NY, USA, 1977.
- 7. J. J. Rotman, An Introduction to Homological Algebra, Academic Press, NewYork, 2009.
- R. Resco, Radicals of Finite Normalizing Extensions, Comm. Algebra, 9(7), (1981), 713-725.
- 9. K. Shimomoto, F-coherent Rings with Applications to Tight Closure Theory, J. Algebra, 338, (2011), 24–34.
- R. Sharp, N. Nossem, Ideals in a Perfect Closure, Linear Growth of Primary Decompositions, and Tight Closure, Transactions Amer. Math. Soc., 356(9), (2004), 3687–3720.
- 11. C. A. Weibel, *An Introduction to Homological Algebra*, vol. 38 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, UK, 1994.
- 12. W. Xue, On Almost Excellent Extensions, Algebra Colloq, 3, (1996), 125-134.

[Downloaded from ijmsi.com on 2025-10-16]

- W. Xue, Rings with Morita Duality, Lect. Notes Math. 1523, Springer-Verlag, Berlin, 1992.
- W. Xue, On n-Presented Modules and Almost Excellent Extensions, J. Algebra, 27(3), (1999), 1091-1102.
- 15. Z. Zhu, n-Cocoherent Rings, n-Cosemihereditary Rings and n-v-Rings, Bull. Iranian Math. Soc., ${\bf 40}(4)$, (2014), 809-822.
- 16. J. Zhan, Z. Tan, Finitely Copresented and Cogenerated Dimensions , $Indian\ J.\ pure\ appl.\ Math.\ {\bf 35}(6),\ (2004),\ 771-781.$
- 17. Z. M. Zhu, J. L. Chen, FCP-Projective Modules and some Rings, $Zhejian\ Univ.\ Sci.\ Ed.,\ 37(2),\ (2010),\ 126-130.$
- D. Zhou, On n-Coherent Rings and (n, d)-Rings, Comm. Algebra, 32(6), (2004), 2425-2441
- D. Zhou, Z. Gong, On Presented Dimensions of Modules and Rings, Int. J. Math. Math. Sci., 2010, Art. ID 256267, 13 pp.