

A New Notions of Quotient Mappings and Networks Through P - I -Convergence

Carlos Granados^{a*}, Laxmi Rathour^b

^aI.E.D. Camilo Torres Tenorio, Barranquilla, Colombia

^bDepartment of Mathematics, National Institute of Technology, Chaltlang,
Aizawl 796 012, Mizoram, India

E-mail: carlosgranadosortiz@outlook.es

E-mail: rathourlaxmi562@gmail.com

ABSTRACT. In this paper, we procure the concepts of P - I -quotient mappings, P - I - cs -networks, P - I - cs' -networks and P - wcs' -networks, and prove some properties of P - I -quotient mappings and P - I - cs' networks, especially P - J -quotient mappings and P - J - cs -networks on an ideal J of $\mathbb{N}$. With these notions, we obtain that if X is a P - J - FU space with a point-countable P - J - cs' -network, then X is a meta-Lindelof space.

Keywords: Ideal convergence, P - I -sequential neighbourhood, P - I -quotient mappings, P - I - cs -networks, P - I - FU spaces.

2000 Mathematics subject classification: 54A20, 40A05.

1. INTRODUCTION

The notion of statistical convergence was originally introduced by H. Fast [10] and H. Steinhaus [24], which is a generalization of the usual notion of convergence. It is doubtless that the study of statistical convergence and its various generalizations has become an active research area in the last few years (see [9, 13, 25]). In particular, P. Kostyrko, T. Salat and W. Wilczynski [12] introduced two interesting generalizations of statistical convergence by using

*Corresponding Author

Received 25 June 2022; Accepted 17 June 2024

©2026 Academic Center for Education, Culture and Research TMU

the notion of ideals of subsets of positive integers, which were named as I and I^* -convergence, and prove some properties of I and I^* -convergence in metric spaces. Moreover, B.K. Lahiri and P. Das [14] discussed I and I^* -convergence in topological spaces. Some further results connected with I and I^* -convergence can be found in [6, 7, 8]. It is well-known that mappings and networks are important tools of investigating topological spaces. Continuous mappings, quotient mappings, pseudo-open mappings, cs -networks, sn -networks, k -networks and so on are the most important tools for studying convergence, sequential spaces, Frechet-Urysohn spaces [16] and generalized metric spaces. Recently, I -quotient mappings and I - cs' -networks was defined in the topological spaces [28] taking into account notions defined in [26] and [27].

Taking into account mentioned above, this paper draws into P - I -quotient mappings and P - I - cs' -networks for an ideal I on $\mathbb{N}$ by using the notion of P - I -convergence defined by Granados [4]. Besides, we discuss about some of their properties.

Throughout this paper, the letter X always denotes a topological space. The cardinality of a set C is denoted by $|C|$. The set of all positive integers, the first infinite ordinal, and the first uncountable ordinal are denoted by $\mathbb{N}$, ω and ω_1 , respectively.

2. PRELIMINARIES

The notion of I -convergence of sequences in a topological space is a generalization of statistical convergence which is based on the ideal of subsets of the set $\mathbb{N}$ of all positive integers. Let $A = 2^{\mathbb{N}}$ be the family of all subsets of $\mathbb{N}$. An ideal $I \subset A$ is a hereditary family of subsets of $\mathbb{N}$ which is stable under finite unions [12]. An ideal I is said to be non-trivial if $I \neq \emptyset$ and $\mathbb{N} \notin I$. A non-trivial ideal $I \subset A$ is called admissible if $\{\{x\} : x \in \mathbb{N}\} \subset I$. It is well-known that every non-trivial ideal defines a dual filter $F_I = \{A \subset \mathbb{N} : \mathbb{N} - A \in I\}$ on $\mathbb{N}$.

EXAMPLE 2.1. The following are admissible ideals on $\mathbb{N}$.

- (1) Let I_f be the family of all finite subsets of $\mathbb{N}$. Then, I_f is an admissible ideal on $\mathbb{N}$.
- (2) Let I_δ be the family of subsets $A \subset \mathbb{N}$ with $\delta(A) = 0$. Then, I_δ is an admissible ideal on $\mathbb{N}$ and the dual filter $F_{I_\delta} = \{A \subset \mathbb{N} : \delta(A) = 1\}$. Here, $\delta(A)$ denotes the asymptotic density of A which is denoted by

$$\delta(A) = \lim_{n \rightarrow \infty} \frac{1}{n} |\{w \in A : w \leq n\}|, \text{ if the limits exists.}$$

Definition 2.2. [4] A sequence $\{x_n\}_{n \in \mathbb{N}}$ in a topological space X is said to be P - I -convergent to a point $x \in X$ provided for any pre-open set U containing x , we have $\{n \in \mathbb{N} : x_n \notin U\} \in I$, which is denoted by P - I - $\lim_{n \rightarrow \infty} x_n = x$ or $x_n \xrightarrow{PI} x$, and the point x is called the P - I -limit of the sequence $\{x_n\}_{n \in \mathbb{N}}$.

Definition 2.3. [5] Let I be an ideal on $\mathbb{N}$ and X be a topological space. Then,

- (1) A subset $F \subset X$ is said to be P - I -closed if for each sequence $\{x_n\}_{n \in \mathbb{N}} \subset F$ with $x_n \xrightarrow{PI} x \in X$, we have $x \in F$.
- (2) A subset $U \subset X$ is said to be P - I -open if $X - U$ is P - I -closed.
- (3) X is called a P - I -sequential space if each P - I -closed subset of X is closed.

Remark 2.4. Every open set is pre-open.

Remark 2.5. Each sequential space is a P - I -sequential space [5].

Definition 2.6. [5] Let I be an ideal on $\mathbb{N}$, X and Y be topological spaces and $f : X \rightarrow Y$ be a mapping.

- (1) f is called preserving P - I -convergence provided for each sequence $\{x_n\}_{n \in \mathbb{N}}$ in X with $x_n \xrightarrow{PI} x$, the sequence $\{f(x_n)\}_{n \in \mathbb{N}}$ P - I -converges to $f(x)$.
- (2) f is called P - I -continuous provided U is P - I -open in Y , then $f^{-1}(U)$ is P - I -open in X .

Remark 2.7. A mapping $f : X \rightarrow Y$ is P - I -continuous if and only if whenever F is P - I -closed in Y , then $f^{-1}(F)$ is P - I -closed in X [5].

Lemma 2.8. [5] Let I be an ideal on $\mathbb{N}$ and X be a topological space. If a sequence $\{x_n\}_{n \in \mathbb{N}}$ P - I -converges to a point $x \in X$ and $\{y_n\}_{n \in \mathbb{N}}$ is a sequence in X with $\{n \in \mathbb{N} : x_n \neq y_n\} \in I$, then the sequence $\{y_n\}_{n \in \mathbb{N}}$ P - I -converges to $x \in X$.

Lemma 2.9. [5] Let I be an ideal on $\mathbb{N}$. The following statements are equivalent for a topological space X and a subset $A \subset X$.

- (1) A is P - I -open.
- (2) $\{n \in \mathbb{N} : x_n \in A\} \notin I$ for each sequence $\{x_n\}_{n \in \mathbb{N}}$ in X with $x_n \xrightarrow{PI} x \in A$.
- (3) $|\{n \in \mathbb{N} : x_n \in A\}| = \omega$ for each sequence $\{x_n\}_{n \in \mathbb{N}}$ in X with $x_n \xrightarrow{PI} x \in A$.

Lemma 2.10. [5] Let X and Y be topological space and $f : X \rightarrow Y$ be a mapping, Then, the following statements hold:

- (1) If f is continuous, then f preserves P - I -convergence.
- (2) If f preserves P - I -convergence, then f is P - I -continuous.

Definition 2.11. [5] Let $A \subset X$ and $\{x_n\}_{n \in \mathbb{N}}$ be a sequence in X . If I is an ideal on $\mathbb{N}$, then $\{x_n\}_{n \in \mathbb{N}}$ is P - I -eventually in A if there is $E \in I$ such that for all $n \in \mathbb{N} - E$, $x_n \in A$.

Remark 2.12. If A is a subset of X with the property that every sequence P - I -converging to a point in A is P - I -eventually in A , then A is P - I -open. When we assume J to be a maximal ideal, the following proposition shows that such sets must coincide with P - J -open sets.

Proposition 2.13. [5] *If J is a maximal ideal of $\mathbb{N}$, then $A \subset X$ is P - J -open if and only if for each P - J -covering sequence $\{x_n\}_{n \in \mathbb{N}}$ with $x_n \xrightarrow{PJ} x \in A$, then $\{x_n\}_{n \in \mathbb{N}}$ is P - J -eventually in A .*

Remark 2.14. By Definition 2.3, the union of a family of P - I -open sets in a topological space is P - I -open. Whenever J is a maximal ideal, the intersection of two P - J -open sets is a P - J -open set.

Proposition 2.15. [5] *If J is a maximal ideal of $\mathbb{N}$ and U, V are two P - J -open subsets of X , then $U \cap V$ is P - J -open in X .*

Remark 2.16. It is well known that the sequential coreflection sX of a space X is the set X endowed with the topology consisting of sequentially open subsets of X . Let J be a maximal ideal of $\mathbb{N}$ and X be a topological space. By Definition 2.3 and Proposition 2.15, the family of all P - J -open subsets of X forms a topology of the set X . The P - J -sequential coreflection of a space X is the set X endowed with the topology consisting of P - J -open subsets of X , which is denoted by P - J - sX . The spaces X and P - J - sX have the same P - J -convergent sequences. Besides, P - J - sX is a P - J -sequential space, further a space X is a P - J -sequential space if and only if P - J - sX .

Throughout this paper, if no otherwise specified, we consider ideal I is always an admissible ideal on $\mathbb{N}$, all mappings are continuous and surjection; and all spaces are Hausdorff.

3. P - I -QUOTIENT MAPPING AND ITS PROPERTIES

In this section, we introduce the notion of P - I -quotient mappings, and obtain some characterizations of P - I -quotient mappings, especially P - J -quotient mappings under a maximal ideal of $\mathbb{N}$.

Definition 3.1. Let I be an ideal on $\mathbb{N}$ and $f : X \rightarrow Y$ be a mapping. Then,

- (1) f is said to be P - I -quotient mapping (or shortly P - I -quotient) provided $f^{-1}(U)$ is P - I -open in X , then U is P - I -open in Y .
- (2) f is said to be P - I -covering mapping (or shortly P - I -covering) if, whenever $\{y_n\}_{n \in \mathbb{N}}$ is a sequence in Y P - I -converging to y in Y , there exist a sequence $\{x_n\}_{n \in \mathbb{N}}$ of points $x_n \in f^{-1}(y_n)$ for all $n \in \mathbb{N}$ and $x \in f^{-1}(y)$ such that $x_n \xrightarrow{PI} x$.

Theorem 3.2. *Every P - I -covering mapping is P - I -quotient.*

Proof. Let X and Y be arbitrary topological spaces, and $f : X \rightarrow Y$ be a P - I -covering mapping. Suppose that H is a non- P - I -closed in Y . Then, there exists a sequence $\{y_n\}_{n \in \mathbb{N}} \subset H$ such that $y_n \xrightarrow{PI} y \notin H$. Since f is P - I -covering, there exists a sequence $\{x_n\}_{n \in \mathbb{N}}$ of points $x_n \in f^{-1}(y_n)$ for all $n \in \mathbb{N}$ and $x \in f^{-1}(y)$ such that $y_n \xrightarrow{PI} y$. We can see that $\{x_n\}_{n \in \mathbb{N}} \subset f^{-1}(H)$

and that $x \notin f^{-1}(H)$. Therefore, $f^{-1}(H)$ is non- P - I -closed. Hence, f is P - I -quotient. $\square$

Lemma 3.3. *The P - J -sequential coreflection P - J - sX is P - J -sequence space and the identity $id_X : P - J - sX \rightarrow X$ is a pre-continuous and P - J -covering mapping.*

Proof. Suppose that U is a pre-open subset in X . Since U is P - J -open in X , it is open in P - J - sX , so $id_X : P - J - sX \rightarrow X$ is pre-continuous. Now, suppose that $\{x_n\}_{n \in \mathbb{N}}$ is a sequence in X which P - J -converges to a point $x \in X$. If V is a pre-open subset in P - J - sX with $x \in V$, the set V is P - J -open in X . By Proposition 2.13, there is $W \in J$ such that for all $n \in \mathbb{N} - W$, $x_n \in V$, i.e., $\{n \in \mathbb{N} : x_n \notin V\} \subset W$ and thus $\{n \in \mathbb{N} : x_n \notin V\} \in J$. Hence, the sequence $x_n \xrightarrow{PJ} x$ in P - J - sX . This means that the spaces X and P - J - sX have the same P - J -convergent sequences. Therefore, $id_X : P - J - sX \rightarrow X$ is a P - J -covering mapping. For each $A \subset X$, by Proposition 2.13, A is P - J -open in pre- J - sX if and only if A is P - I -open in X if and only if A is pre-open in pre- J - sX . Therefore, P - J - sX is a P - J -sequence space. $\square$

Definition 3.4. Let I be an ideal on $\mathbb{N}$, X be a topological space and $P \subset X$. P is called a P - I -sequential neighbourhood of x , if for each sequence $\{x_n\}_{n \in \mathbb{N}}$ P - I -converges to a point $x \in P$, then $\{x_n\}_{n \in \mathbb{N}}$ is P - I -eventually in P , i.e., there is $\tilde{I} \in I$ such that $\{n \in \mathbb{N} : x_n \notin P\} = \tilde{I}$.

Remark 3.5. Let J be a maximal ideal of $\mathbb{N}$ and $A \subset X$. By Proposition 2.13, A is P - J -open in X if and only if A is P - J -sequential neighbourhood of x for each $x \in A$.

Proposition 3.6. *Let J be a maximal ideal of $\mathbb{N}$ and $A \subset X$. If A is not a P - J -sequential neighbourhood of x , then there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in $X - A$ such that $x_n \xrightarrow{PJ} x$.*

Proof. If A is not a P - J -sequential neighbourhood of x , then there is a sequence $\{y_n\}_{n \in \mathbb{N}}$ in X such that $y_n \xrightarrow{PJ} y$, but $\{n \in \mathbb{N} : y_n \notin A\} \notin J$. Since J is a maximal ideal of $\mathbb{N}$, this means that $\{n \in \mathbb{N} : y_n \in A\} \in J$. Now, let $\{n \in \mathbb{N} : y_n \in A\} = \tilde{J} \in J$. And since J is a non-trivial ideal, it follows that $A \neq X$. Taking a point $a \in X - A$. Define a sequence $\{x_n\}_{n \in \mathbb{N}}$ by $x_n = a$ if $n \in \tilde{J}$; $x_n = y_n$ if $n \in \mathbb{N} - \tilde{J}$. Then, the sequence $\{x_n\}_{n \in \mathbb{N}}$ in $X - A$ and $x_n \xrightarrow{PJ} x$ from Lemma 2.9. $\square$

Theorem 3.7. *Let I be an ideal on $\mathbb{N}$. If $f : X \rightarrow Y$ is a P - I -quotient mapping, then for each P - I -convergent sequence $\{y_n\}_{n \in \mathbb{N}}$ in Y with $y_n \xrightarrow{PI} y$, there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X such that $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}})$ and $x_i \xrightarrow{PJ} x \notin f^{-1}(\{y_n\}_{n \in \mathbb{N}})$.*

Proof. Suppose that $f : X \rightarrow Y$ is a $P-I$ -quotient mapping and $\{y_n\}_{n \in \mathbb{N}}$ is sequence in Y with $y_n \xrightarrow{PJ} y$. Without loss of generality, we can assume that $y_n \neq y$ for each $n \in \mathbb{N}$. Now, let $U = Y - \{y_n\}_{n \in \mathbb{N}}$. Then, U is not pre- I -open in Y . Since f is a $P-I$ -quotient mapping, $f^{-1}(U) = f^{-1}(Y - \{y_n\}_{n \in \mathbb{N}}) = X - f^{-1}(\{y_n\}_{n \in \mathbb{N}})$ is not $P-I$ -open in X . Therefore, there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in $X - f^{-1}(U) = f^{-1}(\{y_n\}_{n \in \mathbb{N}})$ such that $x_i \xrightarrow{PJ} x \notin f^{-1}(\{y_n\}_{n \in \mathbb{N}})$. $\square$

In the following theorem we discuss about some properties by using quotient mappings and $P-I$ -quotient mappings.

Theorem 3.8. *Let $f : X \rightarrow Y$ be a mapping. Then, the following statements hold:*

- (1) *If X is a $P-I$ -sequential space and f is quotient, then Y is a $P-I$ -sequential space and f is $P-I$ -quotient.*
- (2) *If Y is a $P-I$ -sequential space and f is $P-I$ -quotient, then f is quotient.*
- (3) *X is a $P-I$ -sequential space if and only if for an arbitrary topological space Y , if f is quotient, then f is $P-I$ -quotient.*

Proof. (1) Let X be a $P-I$ -sequential space and f be quotient. First, we will show that the space Y is a $P-I$ -sequential space, i.e., each pre- I -open set in Y is open. Now, assume that U is pre- I -open in Y . Since f is a quotient mapping and X is a $P-I$ -sequential space, it suffices to show that $f^{-1}(U)$ is pre- I -open in X . Taking an arbitrary arbitrary sequence $\{x_n\}_{n \in \mathbb{N}} \subset X$ such that $x_n \xrightarrow{PI} x \in f^{-1}(U)$ in X . Since f is a quotient mapping from X onto Y , it follows that $f(x_n) \xrightarrow{PI} f(x) \in U$ is pre- I -open in Y , it follows Lemma 2.9 that $|\{n \in \mathbb{N} : f(x_n) \in U\}| = \omega$, i.e., $|\{n \in \mathbb{N} : x_n \in f^{-1}(U)\}| = \omega$. Therefore, $f^{-1}(U)$ is pre- I -open in X . Next, suppose that $U \subset Y$ such that $f^{-1}(U)$ is pre- I -open in X . Then, $f^{-1}(U)$ is open in X since X is a $P-I$ -sequential space, due to f is quotient, U is open in Y , furthermore U is pre- I -open in Y . Therefore, f is $P-I$ -quotient.

(2) Let Y be a $P-I$ -sequential space and f be $P-I$ -quotient. If $f^{-1}(U)$ is open in X , then $f^{-1}(U)$ is pre- I -open in X . Since f is $P-I$ -quotient, U is $P-I$ -open in Y . We can see that Y is a $P-I$ -sequential space, hence U is open in Y . Therefore, f is quotient.

(3) By part (1) and (2) of this theorem, the necessity is obvious.

Sufficiency: If X is not a $P-I$ -sequential space, then there exists a pre- I -closed subset H in X such that H is not closed in X . Now, Let $Y = \{0, 1\}$, define a mapping $f : X \rightarrow Y$ by $f(x) = 0$ if $x \in H$; $f(x) = 1$, if $x \in X - H$. The topology on Y is endowed the continuous quotient topology induced by f . Then, f is quotient. We can check that $f^{-1}(\{1\})$ is not open in X , therefore $\{1\}$ is not open in Y . It follows that the constant sequence $0, 0, \dots, 0, \dots$ $P-I$ -converges to 1 in Y , and hence $\{0\}$ is not pre- I -closed in Y . But $f^{-1}(\{0\}) = H$ is pre- I -closed in X . Therefore, f is not $P-I$ -quotient. $\square$

The following example shows that there exists a continuous and P - I -covering mapping which is not quotient.

EXAMPLE 3.9. Let the space $X = [0, \omega_1]$ be endowed the following topology: the only non-isolated point ω_1 has the semi-neighbourhoods of the usual ordered topology. Then X is not discrete, and there is not any non-trivial convergent sequence in X . The set $Z = [0, \omega_1]$ is endowed the discrete topology. Taking $h : Z \rightarrow X$ is the identity mapping. Obviously, g is continuous, but g is not quotient. Now, suppose that a sequence $\{x_n\}_{n \in \mathbb{N}}$ P - I -converges to x in X . Now, we will show that $\{n \in \mathbb{N} : x_n \neq x\} \in I$, i.e., $\{n \in \mathbb{N} : x_n \notin \{x\}\} \in I$. Let $\{x_n\}_{n \in \mathbb{N}}$ be a sequence in X and $x \in X$, there exists a pre-open set U_x containing x in X with $x_n \notin U_x$ for each $x_n \neq x$. If the sequence $x_n \xrightarrow{PI} x \in X$, then $\{n \in \mathbb{N} : x_n \neq x\} = \{n \in \mathbb{N} : x_n \notin U_x\} \in I$. If the sequence $\{x_n\}_{n \in \mathbb{N}}$ satisfies $\{n \in \mathbb{N} : x_n \neq x\} \in I$ and U is a pre open set containing x in X , we can assume $U_x \subset U$, then $\{n \in \mathbb{N} : x_n \notin U\} \subset \{n \in \mathbb{N} : x_n \notin U_x\} = \{n \in \mathbb{N} : x_n \neq x\} \in I$. Therefore, $x_n \xrightarrow{PI} x$ in Z . Hence, h is P - I -covering.

Theorem 3.10. Let J be a maximal ideal of $\mathbb{N}$ and X be a topological space. Then, X is a P - J -sequential space if and only if each P - J -quotient mapping onto X is quotient.

Proof. The necessity is obtained by Theorem 3.8(2). On the other hand, suppose that every P - J -quotient mapping onto X is quotient. By Lemma 3.3 and Theorem 3.2, the identity $id_X : P - J - sX \rightarrow X$ is a continuous and P - J -quotient mapping, thus it is a quotient mapping. Since P - J - sX is a P - J -sequence space, it follows from Theorem 3.8(1) that X is a P - J -sequential space. $\square$

The following example shows that there exists a quotient mapping which is not P - I -quotient.

EXAMPLE 3.11. The space $X = [0, \omega_1]$ is defined by Example 3.9. The set $Y = \{0, 1\}$ is endowed the topology $\{\emptyset, \{0\}, Y\}$. Define a mapping $f : X \rightarrow Y$ by $f([0, \omega_1]) = \{0\}$, and $f(\omega_1) = 1$. Then, f is a quotient mapping. On the other hand, by Example 3.9, the set $f^{-1}(\{0\}) = [0, \omega_1]$ is pre- I -closed in X . Since the constant sequence $0, 0, \dots, 0, \dots$ P - I -converges to 1 in Y , the set $\{0\}$ is not pre- I -closed in Y . Therefore, f is not P - I -quotient.

Definition 3.12. Let J be a maximal ideal of $\mathbb{N}$ and $A \subset X$. Denote,

- (1) $[A]_{J_P} = \{x \in X : \text{there is a sequence } \{x_n\}_{n \in \mathbb{N}} \text{ in } A \text{ such that } x_n \xrightarrow{PJ} x\}.$
- (2) $(A)_{J_P} = \{x \in X : A \text{ is a } P\text{-}J\text{-sequential neighbourhood of } x\}.$

Definition 3.13. Let J be a maximal ideal of $\mathbb{N}$ and $A \subset X$. A subset $U \subset X$ is said to be P - J -sequential neighbourhood of A if $A \subset (U)_{J_P}$.

Proposition 3.14. *Let J be a maximal ideal of $\mathbb{N}$ and $A \subset X$. Then, $[A]_{J_P} = X - (X - A)_{J_P}$.*

Proof. Suppose that $x \in [A]_{J_P}$, then there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in A such that $x_n \xrightarrow{PJ} x$. Thus, $X - A$ is not a P - J -sequential neighbourhood of x in X . In indeed, if $X - A$ is a P - J -sequential neighbourhood of x in X , then $\{x_n\}_{n \in \mathbb{N}}$ is P - J -eventually in $X - A$, i.e., there is $E \in J$ such that for all $n \in \mathbb{N} - E$, $x_n \in X - A$. Since J is an admissible ideal, this contradicts to $\{x_n\}_{n \in \mathbb{N}}$ in A . Therefore, $x \notin (X - A)_{J_P}$ and further $x \in X - (X - A)_{J_P}$.

On the other hand, assume that $x \in X - (X - A)_{J_P}$, then $x \notin (X - A)_{J_P}$, and hence $X - A$ is not a P - J -sequential neighbourhood of $x \in X$. By Proposition 3.6, there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in A such that $x_n \xrightarrow{PJ} x$. Therefore, $x \in [A]_{J_P}$. $\square$

By Definition 2.3, Definition 3.12 and Proposition 3.14, the following proposition holds.

Proposition 3.15. *Let J be a maximal ideal of $\mathbb{N}$ and $A, B \subset X$. Then,*

- (1) $[\emptyset]_{J_P} = \emptyset$, $\text{Int}(A) \subset (A)_{J_P} \subset A \subset [A]_{J_P} \subset \text{Cl}(A)$, where $\text{Int}(A)$ and $\text{Cl}(A)$ denote interior and closure of A , respectively.
- (2) A is pre- J -open in X if and only if $A = (A)_{J_P}$.
- (3) A is pre- J -closed in X if and only if $A = [A]_{J_P}$.
- (4) If $B \subset A$, then $(B)_{J_P} \subset (A)_{J_P}$ and $[B]_{J_P} \subset [A]_{J_P}$.
- (5) $(A \cap B)_{J_P} = (A)_{J_P} \cap (B)_{J_P}$ and $[A \cup B]_{J_P} = [A]_{J_P} \cup [B]_{J_P}$.

Proof. (1)-(4) are followed directly by the definitions, for that reason we only proof (5).

Since $A \cap B \subset A$ and $A \cap B \subset B$, it follows that $(A \cap B)_{J_P} \subset (A)_{J_P} \cap (B)_{J_P}$. On the other hand, assume that $x \in (A)_{J_P} \cap (B)_{J_P}$. Then, for each sequence $\{x_n\}_{n \in \mathbb{N}}$ in X with $x_n \xrightarrow{PJ} x$, there is $E, F \in J$, such that for each $n \in \mathbb{N} - E$, $x_n \in A$ and for each $n \in \mathbb{N} - F$, $x_n \in B$. Since $E \cup F \in J$ and for each $n \in \mathbb{N} - (E \cup F)$, $x_n \in A \cap B$. This means that $A \cap B$ is a P - I -sequential neighbourhood of x in X . Therefore, $x \in (A \cap B)_{J_P}$.

Now, if we replace $X - A$ with A and $X - B$ with B , it follows that $((X - A) \cap (X - B))_{J_P} = (X - A)_{J_P} \cap (X - B)_{J_P}$. Hence, $[A \cup B]_{J_P} = X - (X - (A \cup B))_{J_P} = X - ((X - A) \cap (X - B))_{J_P} = X - ((X - A) \cap (X - B))_{J_P} = (X - (X - A)_{J_P}) \cup (X - (X - B)_{J_P}) = [A]_{J_P} \cup [B]_{J_P}$. $\square$

Theorem 3.16. *Let J be a maximal ideal of $\mathbb{N}$ and $f : X \rightarrow Y$ be a mapping. Then, the following conditions are equivalent.*

- (1) For each P - J -convergent sequence $\{y_n\}_{n \in \mathbb{N}}$ in Y with $y_n \xrightarrow{PJ} y$, there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X with $x_i \xrightarrow{PJ} x \in f^{-1}(y)$ and $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}})$.
- (2) For each $A \subset Y$, it has $f([f^{-1}(A)]_{J_P}) = [A]_{J_P}$.

- (3) If $y \in [A]_{J_P} \subset Y$, then $f^{-1}(y) \cap [f^{-1}(A)]_{J_P} \neq \emptyset$.
- (4) If $y \in [A]_{J_P} \subset Y$, then there is a point $x \in f^{-1}(y)$ such that whenever V is a P - J -sequential neighbourhood of x , $y \in [f(V) \cap A]_{J_P}$.
- (5) If $y \in [A]_{J_P} \subset Y$, then there is a point $x \in f^{-1}(y)$ such that whenever V is a P - J -sequential neighbourhood of x , $f(V) \cap A \neq \emptyset$.
- (6) For each $y \in Y$, if U is a P - J -sequential neighbourhood of $f^{-1}(y)$, then $f(U)$ is a P - J -sequential neighbourhood of y .

Proof. (1) $\Rightarrow$ (2) Suppose that $x \in [f^{-1}(A)]_{J_P}$. Then, there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in $f^{-1}(A)$ such that $x_n \xrightarrow{PJ} x$. Therefore, $\{f(x_n)\}_{n \in \mathbb{N}} \subset A$ and $f(x_n) \xrightarrow{PJ} f(x)$. This means that $f(x) \in [A]_{J_P}$. Hence, $f([f^{-1}(A)]_{J_P}) \subset [A]_{J_P}$.

On the other hand, assume that $y \in [A]_{J_P}$. Then, there is a sequence $\{y_n\}_{n \in \mathbb{N}}$ in A such that $y_n \xrightarrow{PJ} y$. By hypothesis (1), there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X with $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}}) \subset f^{-1}(A)$ and $x_i \xrightarrow{PJ} x \in f^{-1}(y)$. Thus, $x \in [f^{-1}(A)]_{J_P}$, therefore $y = f(x) \in f([f^{-1}(A)]_{J_P})$, moreover $[A]_{J_P} \subset f([f^{-1}(A)]_{J_P})$.

(2) $\Rightarrow$ (3) Let $y \in [A]_{J_P}$ for each $A \subset Y$. By hypothesis (2), it follows that $y \in f([f^{-1}(A)]_{J_P})$. Therefore, $f^{-1}(y) \cap [f^{-1}(A)]_{J_P} \neq \emptyset$.

(3) $\Rightarrow$ (4) Let $y \in [A]_{J_P} \subset Y$. By hypothesis (3), assume that $x \in f^{-1}(y) \cap [f^{-1}(A)]_{J_P}$. Then, there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ in $f^{-1}(A)$ such that $x_n \xrightarrow{PJ} x$. If V is a P - J -sequential neighbourhood of x , then there is $E \in J$ such that $x_n \in V$ for all $n \in \mathbb{N} - E$. Therefore, $f(x_n) \in f(V) \cap A$ for all $n \in \mathbb{N} - E$ and $f(x_n) \xrightarrow{PJ} f(x)$. Taking a point $a \in f(V) \cap A$. Define a sequence $\{y_n\}_{n \in \mathbb{N}}$ by $y_n = f(x_n)$ if $n \in \mathbb{N} - E$; $y_n = a$ if $n \in E$. Then, $\{y_n\}_{n \in \mathbb{N}} \subset f(V) \cap A$ and $y_n \xrightarrow{PJ} f(x) = y$ from Lemma 2.8. Hence, $y \in [f(V) \cap A]_{J_P}$.

(4) $\Rightarrow$ (5) It is obvious.

(5) $\Rightarrow$ (6) Let $y \in Y$ and U be a P - J -sequential neighbourhood of $f^{-1}(y)$. If $f(U)$ is not a P - J -sequential neighbourhood of y , then $y \in Y - (f(U))_{J_P} = [Y - f(U)]_{J_P}$. By hypothesis (5), it follows that $f(U) \cap (Y - f(U)) = \emptyset$, which is a contradiction.

(6) $\Rightarrow$ (3) Let $y \in [A]_{J_P} \subset Y$. Now, suppose that $f^{-1}(y) \cap [f^{-1}(A)]_{J_P} = \emptyset$. Then, $f^{-1}(y) \subset X - [f^{-1}(A)]_{J_P} = (X - f^{-1}(A))_{J_P}$. This means that $X - f^{-1}(A)$ is a P - J -sequential neighbourhood of $f^{-1}(y)$. By hypothesis (6), $y \in (f(X - f^{-1}(A)))_{J_P} = (Y - A)_{J_P} = Y - [A]_{J_P}$, which is a contradiction.

(3) $\Rightarrow$ (1) Let $\{y_n\}_{n \in \mathbb{N}}$ be a P - J -convergent sequence in Y with $y_n \xrightarrow{PJ} y$. Now, put $A = \{y_n\}_{n \in \mathbb{N}}$, then $A \in [A]_{J_P}$. By hypothesis (3), there is $x \in f^{-1}(y) \cap [f^{-1}(A)]_{J_P}$. Therefore, there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X with $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(A) \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}})$ and $x_i \xrightarrow{PJ} x \in f^{-1}(y)$. $\square$

Remark 3.17. One of the above six conditions can deduce that f is a P - J -quotient mapping. In indeed, let U be non- P - I -closed in Y . Then, there is a

sequence $\{y_n\}_{n \in \mathbb{N}}$ in U P - J -converging to $y \in Y - U$. Thus $y \neq y_n$ for each $n \in \mathbb{N}$. By the assumption of the condition (1), there is a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X such that $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}}) \subset f^{-1}(U)$ and $x_i \xrightarrow{PJ} x \in f^{-1}(y) \notin f^{-1}(U)$. This implies that $f^{-1}(U)$ is non- P - J -closed in X . Therefore, f is a P - J -quotient mapping.

Remark 3.18. If the maximal ideal J is replaced by I_f in Theorem 3.16, then (1) $\Leftrightarrow$ (2) $\Leftrightarrow$ (3) $\Leftrightarrow$ (4) $\Leftrightarrow$ (5) $\Leftrightarrow$ (6) $\Leftrightarrow$ f is a P - I_f -quotient mapping. But the following example shows that there exist a T_1 space X , an ideal I of $\mathbb{N}$ and a P - I -quotient mapping f such that f does not satisfy the condition (6) of Theorem 3.16.

EXAMPLE 3.19. Let $I = \{A \subset \mathbb{N} : A \text{ contains at most only finite odd positive integers}\}$. Then, I is an admissible ideal of $\mathbb{N}$. Now, Let Y be the set ω which is endowed with the finite complement topology. Then, Y is a first-countable T_1 -space. Now, putting $X_0 = Y - \{0\}$ and $X_1 = \{2k : k \in \omega\}$ as the subspaces of the space Y and $X = X_0 \oplus X_1$. A mapping $f : X \rightarrow Y$ is defined by the natural mapping. It can see that the mapping f is a continuous quotient mapping. Since X_0 and X_1 are first-countable space, X is a first-countable space. Hence, X is a P - I -sequential space. By Theorem 3.8, it follows that f is a P - I -quotient mapping.

Next, note that the set X_1 is open in X and $f^{-1}(0) \subset X_1$ and hence X_1 is a P - I -sequential neighbourhood of $f^{-1}(0)$. For each open neighbourhood U of 0 in Y , $\{n \in \mathbb{N} : n \notin U\}$ is a finite subset, therefore $\{n \in \mathbb{N} : n \notin U\} \in I$. This means that the sequence $\{n\}_{n \in \mathbb{N}}$ in Y satisfies $n \xrightarrow{PI} 0 \in f^{-1}(y)$. But, $\{n \in \mathbb{N} : n \notin f(X_1)\} = \{2k+1, k \in \omega\} \notin I$. Thus $f(X_1)$ is not a P - I -sequential neighbourhood of 0 in Y .

Remark 3.20. Theorem 3.7 is different from Theorem 3.16(1). In Theorem 3.16(1), $x_i \xrightarrow{PJ} x \in f^{-1}(y)$. But we do not know whether the P - I -limit point x in $f^{-1}(y)$ or not in Theorem 3.7.

4. ON SPACES WITH P - I -cs-NETWORKS, P - I -cs'-NETWORKS AND P -wcs'-NETWORKS

In this section, we introduce the notions of P - I -cs-networks, P - I -cs'-networks and P -wcs'-networks for a space X ; and obtain that if X is an P - J -FU space with a point-countable P - J -cs'-network, then X is a meta-Lindelof space, for an ideal J of $\mathbb{N}$.

Definition 4.1. Let I be an ideal on $\mathbb{N}$, X be a topological space and P be a cover of X .

- (1) [15] P is a network of X if whenever $x \in U$ with U open in X , then $x \in \tilde{P} \subset U$ for some $\tilde{P} \in P$.

- (2) P is a P -network of X if whenever $x \in U$ with U pre-open in X , then $x \in \tilde{P} \subset U$ for some $\tilde{P} \in P$.
- (3) P is called P - I -cs-network of X if whenever $\{x_n\}_{n \in \mathbb{N}}$ is a sequence in X P - I -covering to a point $x \in U$ with U pre-open in X , then $\{x_n\}_{n \in \mathbb{N}}$ is P - I -eventually in $\tilde{P}$ and $x \in \tilde{P} \subset U$ for some $\tilde{P} \in P$.
- (4) P is called P - I -cs'-network of X if whenever $\{x_n\}_{n \in \mathbb{N}}$ is a sequence in X P - I -covering to a point $x \in U$ with U pre-open in X , then there is $\tilde{P} \in P$ and some $n_0 \in \mathbb{N}$ such that $\{x, x_{n_0}\} \subset \tilde{P} \subset U$.
- (5) P is called P - I -wcs'-network of X if whenever $\{x_n\}_{n \in \mathbb{N}}$ is a sequence in X P - I -covering to a point $x \in U$ with U pre-open in X , then there is $\tilde{P} \in P$ and some $n_0 \in \mathbb{N}$ such that $\{x_{n_0}\} \subset \tilde{P} \subset U$.

By Definition 4.1, we have the following diagram:

$$P\text{-}I\text{-}cs\text{-}networks \implies P\text{-}I\text{-}cs'\text{-}networks \implies P\text{-}I\text{-}wcs'\text{-}networks \implies P\text{-}networks.$$

Proposition 4.2. *Every P -network is a network.*

Proof. Let P be a S -network and U be an open subset of X containing x . Then, U is pre-open in X . Since P is a P -network for X , there is some $\tilde{P} \in P$ such that $x \in \tilde{P} \subset U$. Thus P is a network for X . $\square$

Definition 4.3. Let J be a maximal ideal of $\mathbb{N}$ and X be a topological space. U is said to be P - J -sn-cover of X , if $\{(\tilde{U})_{J_P} : \tilde{U} \in U\}$ is a cover of X .

Theorem 4.4. *Each P - I -cs-network is preserved by a P - I -covering mapping.*

Proof. Let $f : X \rightarrow Y$ be a P - I -covering mapping and P be a P - I -cs-network of X . Now, suppose that $\{y_n\}_{n \in \mathbb{N}}$ is a sequence P - I -converging to a point $y \in U$ with U pre-open in Y . Since f is a P - I -covering mapping, there exist a sequence of points $x_n \in f^{-1}(y_n)$ for all $n \in \mathbb{N}$ and $x \in f^{-1}(y)$ such that $x_n \xrightarrow{PI} x$. Since P is a P - I -cs-network of X , there is some $\tilde{P} \in P$ such that $\{x_n\}_{n \in \mathbb{N}}$ is P - I -eventually in $\tilde{P}$ and $x \in \tilde{P} \subset f^{-1}(U)$. Thus there is $E \in I$ such that $\{n \in \mathbb{N} : x_n \notin \tilde{P}\} \subset E$. Note that $\{n \in \mathbb{N} : y_n \notin f(\tilde{P})\} \subset \{n \in \mathbb{N} : x_n \notin \tilde{P}\} \subset E$, hence $y_n \in f(\tilde{P})$ for all $n \in \mathbb{N} - E$, i.e. $\{y_n\}_{n \in \mathbb{N}}$ is P - I -eventually in $f(\tilde{P})$ and $y \in f(\tilde{P}) \subset U$. This means that $f(P) = \{f(\tilde{P}) : \tilde{P} \in P\}$ is a P - I -cs-network of Y . $\square$

Corollary 4.5. *Each P - I -cs'-network is preserved by a P - I -covering mapping.*

Proof. Proof follows from Theorem 4.4. $\square$

Theorem 4.6. *Each P - I -wcs'-network is preserved by a P - I -quotient mapping.*

Proof. Let $f : X \rightarrow Y$ be a P - I -quotient mapping and P be a P - I -wcs'-network of X . Now, suppose that $\{y_n\}_{n \in \mathbb{N}}$ is a sequence P - I -converging to a point $y \in U$ with U pre-open in Y . Since f is a P - I -quotient mapping,

there exist a sequence $\{x_i\}_{i \in \mathbb{N}}$ in X such that $\{x_i\}_{i \in \mathbb{N}} \subset f^{-1}(\{y_n\}_{n \in \mathbb{N}})$ and $x_i \xrightarrow{PI} x \in f^{-1}(\{y_n\}_{n \in \mathbb{N}})$. And because P is a P - I - wcs' -network of X , there is some $\tilde{P}_0 \in P$ and $i_0 \in \mathbb{N}$ such that $\{x_{i_0}\} \subset \tilde{P}_0 \subset f^{-1}(U)$. And hence $\{f(x_{i_0})\} \subset \{y_{n_0}\} \subset f(\tilde{P}_0) \subset U$ for some $n_0 \in \mathbb{N}$. This implies that $f(P) = \{f(\tilde{P}) : \tilde{P} \in P\}$ is a P - I - wcs' -network of Y . $\square$

Lemma 4.7. *Let J be maximal ideal of $\mathbb{N}$ and P be a family of subsets of X . Then, P is a P - J - cs' -network of X if and only if, whenever U is a pre-open set containing x , $\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$ is a P - J -sequential neighbourhood of x .*

Proof. Necessity: Let U be a pre-open set containing x . If $\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$ is not a P - J -sequential neighbourhood of x , then there is a sequence $\{x_n\}_{n \in \mathbb{N}}$ such that $x_n \xrightarrow{PJ} x$ and $x_n \notin \bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$. Since P is a P - J - cs' -network of X , there is $\tilde{P}_0 \in P$ and $n_0 \in \mathbb{N}$ such that $\{x, x_{n_0}\} \subset \tilde{P}_0 \subset U$, which is a contradiction.

Sufficiency: Suppose that $x_n \xrightarrow{PJ} x \in U \in \tau_X$ and $\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$ is a P - J -sequential neighbourhood of x . Then $\{x_n\}_{n \in \mathbb{N}}$ is P - J -eventually in $\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$. Hence there exists $n_0 \in \mathbb{N}$ such that $x_{n_0} \in \bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset U\}$. And hence there is $\tilde{P}_0 \in P$ such that $x_{n_0} \in \tilde{P}_0$ and $x \in \tilde{P}_0 \subset U$. Thus $\{x, x_{n_0}\} \subset \tilde{P}_0 \subset U$. This means that P is a P - J - cs' -network of X . $\square$

Theorem 4.8. *Let J be a maximal ideal of $\mathbb{N}$ and a space X be of a point-countable P - J - cs' -network. Then each open cover of X has a point-countable P - J - sn refinement.*

Proof. Suppose that P is a point-countable P - J - cs' -network for a space X . Now, let $U = \{\tilde{U}_\alpha\}_{\alpha < \gamma}$ be a pre-open cover of X , where γ is an ordinal. For each $\alpha < \gamma$, put

$$V_\alpha = \bigcup\{\tilde{P} \in P : \tilde{P} \subset \tilde{U}_\alpha, \tilde{P} \not\subset \tilde{U}_\beta \text{ if } \beta < \alpha\}.$$

It is clear that $V_\alpha \subset \tilde{U}_\alpha$. Next we shall show that the family $\mathcal{V} = \{V_\alpha\}_{\alpha < \gamma}$ is a point-countable P - J - sn -cover of X . For each $x \in X$, let $\alpha(x) = \min\{\alpha < \gamma : x \in \tilde{U}_\alpha\}$. Then $x \in \tilde{U}_{\alpha(x)}$ and

$$\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset \tilde{U}_{\alpha(x)}\} \subset \bigcup\{\tilde{P} \in P : \tilde{P} \subset \tilde{U}_\alpha, \tilde{P} \not\subset \tilde{U}_\beta \text{ if } \beta < \alpha(x)\}.$$

Since P is an P - J - cs' -network for a space X , it follows from Lemma 4.7 that

$$\begin{aligned} x &\in (\bigcup\{\tilde{P} \in P : x \in \tilde{P} \subset \tilde{U}_{\alpha(x)}\})_{J_P} \\ &\subset (\bigcup\{\tilde{P} \in P : \tilde{P} \subset \tilde{U}_\alpha, \tilde{P} \not\subset \tilde{U}_\beta \text{ if } \beta < \alpha(x)\})_{J_P} \\ &= (V_{\alpha(x)})_{J_P}. \end{aligned}$$

This means that $\mathcal{V} = \{V_\alpha\}_{\alpha < \gamma}$ is a P - J - sn -cover of X . $\square$

Definition 4.9. [5] A space X is called P - I -Frechet-Urysohn (or shortly P - I - FU space) if for each $A \subset X$ and each $x \in Cl(A)$, there exists a sequence in A P - I -converging to the point x in X . Here, $Cl(A)$ denotes closure of A .

Definition 4.10. [15] A space X is called a meta-Lindelof space if each open cover of X has a point-countable open refinement.

Corollary 4.11. Let P be a maximal ideal of $\mathbb{N}$. If X is a P - J - FU space with a point-countable P - J - cs' -network, then X is a meta-Lindelof space.

Proof. X is a P - J - FU if and only if $Cl(A) = [A]_{J_P}$, for each $A \subset X$ if and only if $Int(A) = (A)_{J_P}$ for each $A \subset X$. $\square$

Remark 4.12. In the last corollary, $Int(A)$ denotes interior of A .

Theorem 4.13. Let J be a maximal ideal of $\mathbb{N}$. The following are equivalent for a space X .

- (1) P - J - sX is a S - J -Frechet-Urysohn space.
- (2) $Cl_{P-J-sX}(A) = [A]_{J_P}$, for each $A \subset X$.
- (3) $[A]_{J_P}$ is P - J -closed in X , for each $A \subset X$.
- (4) $(A)_{J_P}$ is P - J -open in X , for each $A \subset X$.

Proof. Since the spaces X and P - J - sX have the same P - J -convergent sequences, by the Definitions 4.9 and 4.10; and Proposition 3.14, it follows that (1) $\Leftrightarrow$ (2) and (3) $\Leftrightarrow$ (4). Hence, it suffices to show that (2) $\Leftrightarrow$ (3). If $Cl_{P-J-sX}(A) = [A]_{J_P}$, then $[A]_{J_P}$ is closed in P - J - sX , and hence $[A]_{J_P}$ is P - J -closed in X , for each $A \subset X$. On the other hand, if $[A]_{J_P}$ is P - J -closed in X , then $[A]_{J_P}$ is closed in P - J - sX , and further $Cl_{P-J-sX}(A) = [A]_{J_P}$, for each $A \subset X$. $\square$

5. CONCLUSION

In this paper we have studied a new kind of quotient and networks properties by using the concept of P - I -open set which was previously defined in [4]. With this paper we propose a new area of research that mix the notions of I -convergence and open sets in topological spaces. We expect that authors can extend these notions in other spaces such that β - I -open sets, α - I -open sets and in the results obtained by [1, 2, 3, 11, 17, 18, 19, 20, 21, 22, 23] can be extended if it is possible.

ACKNOWLEDGMENTS

Thankful to respected Referee for their careful reading of the paper and several valuable suggestions which has improved the presentation of the paper.

REFERENCES

1. L. Deepmala, V. Mishra, L. Mishra, Trigonometric Approximation of Signals (Functions) Belonging to the $W(L_r, \xi(t))$, $(r \geq 1)$ — class by (E, q) $q > 0$ —means of the Conjugate Series of its Fourier Series, *Global Journal of Mathematical Sciences*, **2**(2), (2014), 61-69.
2. A. Gairola, A. Singh, L. Rathour, V. N. Mishra, Improved Rate of Approximation by Modification of Baskakov Operator, *Operators and Matrices*, **16**(4), (2022), 1097-1123.
3. A. Gairola, S. Maindola, L. Rathour, L. Mishra, V. Mishra, Better Uniform Approximation by New Bivariate Bernstein Operators, *International Journal of Analysis and Applications*, **20**, (2022), 1-19.
4. C. Granados, Pre- I_{sn} -open Sets and Some Notions Related to Pre-I-convergence, *Journal of Mathematical Control Science and Applications*, **7**(1), (2021), 1-9.
5. C. Granados, A Note on P-I-convergence, *Divulgaciones Matemáticas*, **21**(1-2), (2020), 1-8.
6. C. Granados, A Generalized of the Strongly Cesáro Ideal Convergence Through Double Sequences Spaces, *International Journal of Applied Mathematics*, **34**(3), (2021), 525-533.
7. C. Granados, J. Bermúdez, I_2 -localized Double Sequences in Metric Spaces, *Advances in Mathematics: Scientific Journal*, **10**(6), (2021), 2877-2885.
8. C. Granados, New Notions of Triple Sequences on Ideal Spaces in Metric Spaces, *Advances in the Theory of Nonlinear Analysis and its Applications*, **5**(3), (2021), 363-368.
9. C. Granados, A. Dhital, Statistical Convergence of Double Sequences in Neutrosophic Normed Spaces, *Neutrosophic Sets and Systems*, **42**, (2021), 333-344.
10. H. Fast, Sur la Convergence Statistique, *Colloquium Mathematicum*, **2**, (1951), 241-244.
11. K. Khatri, A Study on Approximation of Functions in Banach Spaces Using Summability Methods, Ph. D. Thesis, S. V. National Institute of Technology, Surat, 2014.
12. P. Kostyrko, T. Salat, W. Wilczyński, I -convergence, *Real Analysis Exchange*, **26**, (2000/2001), 669-686.
13. V. Kumar, M. Mursaleen, On (λ, μ) -statistical Convergence of Double Sequences on Intuitionistic Fuzzy Normed Spaces, *Filomat*, **25**(2), (2011), 109-120.
14. B. Lahiri, P. Das, I and I^* -convergence in Topological Spaces, *Mathematica Bohemica*, **130**(2), (2005), 153-160.
15. S. Lin, X. Ling, Y. Ge, Point-regular Covers and Sequence-covering Compact Mappings, *Topology and its Applications*, **271**, (2020), 106987.
16. S. Lin, Z. Yun, Generalized Metric Spaces and Mapping, *Atlantis Studies in Mathematics*, **6**, (2016), 1-11.
17. L. Mishra, V. Mishra, K. Khatri, D. Deepmala, On The Trigonometric Approximation of Signals Belonging to Generalized Weighted Lipschitz $W(L^r, \xi(t))$ $(r \geq 1)$ — Class by Matrix $(C^1.N_p)$ Operator of Conjugate Series of its Fourier Series, *Applied Mathematics and Computation*, **237**, (2014), 252-263.
18. L. Mishra, M. Raiz, L. Rathour, V. Mishra, Tauberian Theorems for Weighted Means of Double Sequences in Intuitionistic Fuzzy Normed Spaces, *Yugoslav Journal of Operations Research*, **32**(3), (2022), 377-388.
19. V. Mishra, K. Khatri, L. Mishra, Using Linear Operators to Approximate Signals of Lip (α, p) , $(p \geq 1)$ -Class, *Filomat*, **27**(2), (2013), 353-363.
20. V. Mishra, K. Khatri, L. Mishra, D. Deepmala, Inverse Result in Simultaneous Approximation by Baskakov-Durrmeyer-Stancu Operators, *Journal of Inequalities and Applications*, **2013**, (2013), 586.
21. V. Mishra, Some Problems on Approximations of Functions in Banach Spaces, Ph.D. Thesis, Indian Institute of Technology, India, 2007.

22. V. Mishra, L. Mishra, Trigonometric Approximation of Signals (Functions) in L_p -norm, *International Journal of Contemporary Mathematical Sciences*, **7**(19), (2012), 909-918.
23. L. Mishra, *On Existence and Behavior of Solutions to Some Nonlinear Integral Equations with Applications*, Ph.D. Thesis, National Institute of Technology, India, 2017.
24. H. Steinhaus, Sur la Convergence Ordinaire et la Convergence Asymptotique, *Colloquium Mathematicum*, **2**, (1951), 73-74.
25. N. Subramanian, A. Esi, Wijsman Rough Lacunary Statistical Convergence on I Cesaro Triple Sequences, *International Journal of Analysis and Applications*, **16**(5), (2018), 643-653.
26. X. Zhou, L. Liu, On I -covering Mappings and I -covering Mappings, *Journal of Mathematical Research with Applications*, **40**(1), (2020) 47-56.
27. X. Zhou, L. Liu, S. Lin, On Topological Spaces Defined by I -convergence, *Bulletin of the Iranian Mathematical Society*, **46**, (2020), 675-692.
28. X. Zhou, On I -quotient Mappings and I - cs' -networks Under a Maximal Ideal, *Applied General Topology*, **21**(2), (2020), 235-246.