

## Some Generalizations of Locally Closed Sets

Shyamapada Modak<sup>\*,a</sup> and Takashi Noiri<sup>b</sup>

<sup>a</sup>Department of Mathematics, University of Gour Banga  
P.O. Mokdumpur, Malda 732 103, India.

<sup>b</sup>2949-1 Shiokita-cho, Hinagu, Yatsushiro-shi  
Kumamoto-ken, 869-5142 JAPAN.

E-mail: spmodak2000@yahoo.co.in

E-mail: t.noiri@nifty.com

**ABSTRACT.** Arenas et al. [1] introduced the notion of  $\lambda$ -closed sets as a generalization of locally closed sets. In this paper, we introduce the notions of  $\lambda$ -locally closed sets,  $\Lambda_\lambda$ -closed sets and  $\lambda g$ -closed sets and obtain some decompositions of closed sets and continuity in topological spaces.

**Keywords:**  $\lambda$ -Open set,  $\lambda$ -Locally closed set,  $\Lambda_\lambda$ -Closed set,  $\lambda g$ -Closed set, Decompositions of continuity.

**2000 Mathematics subject classification:** 54A05, 54C08.

### 1. INTRODUCTION AND PRELIMINARIES

The study of locally closed sets was introduced by Bourbaki [3] in 1966 then the authors Ganster and Reilly [6] have studied it extensively. A subset  $A$  of a topological space  $X$  is called locally closed if  $A = U \cap F$ , where  $U$  is open and  $F$  is closed. It is interesting that a locally closed set is a generalization of both open sets and closed sets. The generalization has also been discussed in completely regular Hausdorff spaces [5] and has also been done on algebra with topology in [12] and [2].

---

\*Corresponding Author

Received 23 November 2016; Accepted 29 March 2017

©2019 Academic Center for Education, Culture and Research TMU

In this paper we consider a new type of sets in the topological space which is called  $\lambda$ -open sets. A set is said to be  $\lambda$ -open if it contains a nonempty open set. This idea is not a new idea. In literature, semi-open sets [7] and  $\alpha$ -sets [11] are examples of that type of sets although preopen sets [10] is not an example of it. Because: let  $\mathbf{R}$  be the usual real line and  $Q$  the rational numbers. Then  $\text{Cl}(Q) = \mathbf{R}$  and  $Q \subseteq \text{Int}(\text{Cl}(Q)) = \mathbf{R}$  (where ‘Cl’ and ‘Int’ denote the closure and interior operators, respectively). But  $Q$  does not contain nonempty open set. However Dontchev [4] has introduced an  $S$ -space: A topological space  $X$  is called an  $S$ -space if every subset which contains a non-void open subset is open. But the concept of  $\lambda$ -open sets is different from Dontchev’s  $S$ -spaces.

**Definition 1.1.** A subset  $A$  of a topological space  $X$  is said to be  $\lambda$ -open if  $A$  contains a nonempty open set. The complement of a  $\lambda$ -open set is said to be  $\lambda$ -closed.

For a subset  $A$  of a topological space  $X$ ,  $\text{Int}_\lambda(A)$  and  $\text{Cl}_\lambda(A)$  are defined as follows:

**Definition 1.2.** Let  $X$  be a topological space and  $A$  be a subset of  $X$ .

$$\text{Int}_\lambda(A) = \cup\{U : U \subseteq A, U \text{ is } \lambda\text{-open in } X\};$$

$$\text{Cl}_\lambda(A) = \cap\{F : A \subseteq F, F \text{ is } \lambda\text{-closed in } X\}.$$

**Lemma 1.3.** Let  $X$  be a topological space and  $A, B$  subsets of  $X$ .

- (1) if  $A \subseteq B$ , then  $\text{Int}_\lambda(A) \subseteq \text{Int}_\lambda(B)$  and  $\text{Cl}_\lambda(A) \subseteq \text{Cl}_\lambda(B)$ ,
- (2)  $X \setminus \text{Int}_\lambda(A) = \text{Cl}_\lambda(X \setminus A)$ ,
- (3) For any index set  $\Delta$ , if  $A_\alpha$  is  $\lambda$ -open (resp.  $\lambda$ -closed), then  $\cup\{A_\alpha : \alpha \in \Delta\}$  is  $\lambda$ -open (resp.  $\cap\{A_\alpha : \alpha \in \Delta\}$  is  $\lambda$ -closed),
- (4)  $\text{Int}_\lambda(A)$  is  $\lambda$ -open and  $\text{Cl}_\lambda(A)$  is  $\lambda$ -closed.

**Remark 1.4.** The finite intersection of  $\lambda$ -open sets need not be  $\lambda$ -open. Let  $\mathbf{R}$  be the usual real line,  $A = (-1, 0]$  and  $B = [0, 1)$ . The  $A$  and  $B$  are  $\lambda$ -open but  $A \cap B = \{0\}$  is not  $\lambda$ -open.

We generalize the locally closed set by using  $\lambda$ -open sets.

## 2. $\lambda$ -LOCALLY CLOSED SETS

**Definition 2.1.** A subset  $A$  of a topological space  $X$  is said to be  $\lambda$ -locally closed if  $A = U \cap F$ , where  $U$  is  $\lambda$ -open and  $F$  is closed.

**Corollary 2.2.** Let  $f : X \rightarrow Y$  be a continuous function. If  $L$  is a  $\lambda$ -locally closed subset of  $Y$ , then  $f^{-1}(L)$  is  $\lambda$ -locally closed in  $X$ .

From Definition 1.1 it is obvious that every locally closed set is  $\lambda$ -locally closed. But the converse need not hold in general.

**EXAMPLE 2.3.** Let  $X = \{a, b, c, d\}$ ,  $\tau = \{\emptyset, X, \{a\}\}$ . Then  $C(X)$ (all closed sets in  $X$ ) =  $\{\emptyset, X, \{b, c, d\}\}$ . And  $\lambda$ -open sets are:  $\emptyset$ ,  $X$ ,  $\{a\}$ ,  $\{a, b\}$ ,  $\{a, b, c\}$ ,  $\{a, c\}$ ,

$\{a, d\}$ ,  $\{a, b, d\}$ ,  $\{a, c, d\}$ . Therefore,  $\{d\} = \{a, d\} \cap \{b, c, d\}$  is a  $\lambda$ -locally closed set but it is not a locally closed set in  $X$ .

**Remark 2.4.** A subset  $A$  of a topological space  $X$  is  $\lambda$ -locally closed if and only if  $X \setminus A$  is the union of a  $\lambda$ -closed set and an open set.

**Remark 2.5.** For a subset of a topological space, the following hold:

- (1) Every  $\lambda$ -open set is  $\lambda$ -locally closed,
- (2) Every closed set is  $\lambda$ -locally closed.

**Theorem 2.6.** For a subset  $A$  of a topological space  $X$ , the following are equivalent:

- (1)  $A$  is  $\lambda$ -locally closed;
- (2)  $A = U \cap \text{Cl}(A)$  for some  $\lambda$ -open set  $U$ ;
- (3)  $A \cup (X \setminus \text{Cl}(A))$  is  $\lambda$ -open;
- (4)  $A \subseteq \text{Int}_\lambda[A \cup (X \setminus \text{Cl}(A))]$ ;
- (5)  $\text{Cl}(A) \setminus A$  is  $\lambda$ -closed.

*Proof.* (1)  $\Rightarrow$  (2): Suppose  $A$  is  $\lambda$ -locally closed. Then  $A = U \cap F$  where  $U$  is  $\lambda$ -open and  $F$  is closed. Then  $\text{Cl}(A) = \text{Cl}(U \cap F) \subseteq \text{Cl}(F) = F$ . Then  $A \subseteq U \cap \text{Cl}(A) \subseteq U \cap F = A$  and hence  $A = U \cap \text{Cl}(A)$ .

(2)  $\Rightarrow$  (3):  $X \setminus [A \cup (X \setminus \text{Cl}(A))] = (X \setminus A) \cap \text{Cl}(A) = \text{Cl}(A) \setminus A = \text{Cl}(A) \setminus (U \cap \text{Cl}(A)) = \text{Cl}(A) \setminus U = \text{Cl}(A) \cap (X \setminus U)$ . Since  $U$  is  $\lambda$ -open,  $\text{Cl}(A) \cap (X \setminus U)$  is  $\lambda$ -closed and hence  $A \cup (X \setminus \text{Cl}(A))$  is  $\lambda$ -open.

(3)  $\Rightarrow$  (4): Since  $A \cup (X \setminus \text{Cl}(A))$  is a  $\lambda$ -open set containing  $A$ , it is obvious that  $A \subseteq \text{Int}_\lambda[A \cup (X \setminus \text{Cl}(A))]$ .

(4)  $\Rightarrow$  (1):  $A = A \cap \text{Cl}(A) \subseteq \text{Int}_\lambda[A \cup (X \setminus \text{Cl}(A))] \cap \text{Cl}(A) \subseteq [A \cup (X \setminus \text{Cl}(A))] \cap \text{Cl}(A) = A \cap \text{Cl}(A) = A$ . Therefore,  $A = \text{Int}_\lambda[A \cup (X \setminus \text{Cl}(A))] \cap \text{Cl}(A)$  and  $A$  is  $\lambda$ -locally closed.

(3)  $\Leftrightarrow$  (5): It is obvious.  $\square$

The union of two  $\lambda$ -locally closed sets need not be  $\lambda$ -locally closed.

**EXAMPLE 2.7.** Let  $X = \{a, b, c, d\}$ ,  $\tau = \{\emptyset, X, \{a, b\}, \{c, d\}\}$ . Then  $C(X) = \{\emptyset, X, \{c, d\}, \{a, b\}\}$  and  $\lambda$ -open sets are:  $\emptyset$ ,  $X$ ,  $\{a, b\}$ ,  $\{c, d\}$ ,  $\{a, b, c\}$ ,  $\{a, b, d\}$ ,  $\{a, c, d\}$ ,  $\{b, c, d\}$ .  $\lambda$ -locally closed sets are:  $\emptyset$ ,  $X$ ,  $\{a, b\}$ ,  $\{c, d\}$ ,  $\{a, b, c\}$ ,  $\{a, b, d\}$ ,  $\{a, c, d\}$ ,  $\{b, c, d\}$ ,  $\{c\}$ ,  $\{d\}$ ,  $\{a\}$ ,  $\{b\}$ . Therefore,  $\{a\}$  and  $\{c\}$  are  $\lambda$ -locally closed sets but their union  $\{a, c\}$  is not a  $\lambda$ -locally closed set.

### 3. $\Lambda_\lambda$ -CLOSED SETS

Locally closed sets in a topological space are introduced and investigated in [3] and [6]. As a generalization of locally closed sets, Arenas et al. [1] introduced the notion of  $\lambda$ -closed sets in a topological space. In this section, we introduce the notion of  $\Lambda_\lambda$ -closed sets which is a generalization of  $\lambda$ -closed sets. We obtain some characterizations of  $\Lambda_\lambda$ -closed sets and obtain decompositions of closed sets.

**Definition 3.1.** Let  $X$  be a topological space and  $A$  a subset of  $X$ . The subset  $\Lambda_\lambda(A)$  is defined as follows:  $\Lambda_\lambda(A) = \cap\{U : A \subseteq U, U \text{ is } \lambda\text{-open}\}$ .

A subset  $A$  is called a  $\Lambda_\lambda$ -set if  $A = \Lambda_\lambda(A)$ . If  $U$  is open in Definition 3.1, then a  $\Lambda_\lambda$ -set  $A$  is called a  $\Lambda$ -set [9].

**Lemma 3.2.** For any subsets  $A$  and  $B$  of a topological space  $X$ , the following hold:

- (1)  $A \subseteq \Lambda_\lambda(A)$ ,
- (2) If  $A \subseteq B$ , then  $\Lambda_\lambda(A) \subseteq \Lambda_\lambda(B)$ ,
- (3)  $\Lambda_\lambda(\Lambda_\lambda(A)) = \Lambda_\lambda(A)$ ,
- (4)  $\Lambda_\lambda(\cap_{\alpha \in \Delta} A_\alpha) \subseteq \cap_{\alpha \in \Delta} \Lambda_\lambda(A_\alpha)$  for any index set  $\Delta$ .

**Lemma 3.3.** For any subset  $A$  of a topological space  $X$ , the following hold:

- (1)  $\Lambda_\lambda(A)$  is a  $\Lambda_\lambda$ -set,
- (2) If  $A$  is  $\lambda$ -open, then  $A$  is a  $\Lambda_\lambda$ -set,
- (3) If  $A_\alpha$  is a  $\Lambda_\lambda$ -set for each  $\alpha \in \Delta$ , then  $\cap_{\alpha \in \Delta} A_\alpha$  is a  $\Lambda_\lambda$ -set.

*Remark 3.4.* The converse of Lemma 3.3 (2) need not hold as shown by the following example: Let  $\mathbf{R}$  be the usual real line and  $A = \{0\}$ . Then  $A$  is a  $\Lambda_\lambda$ -set but it is not  $\lambda$ -open. Because  $\{0\} \subseteq \Lambda_\lambda(\{0\}) \subseteq (-1, 0] \cap [0, 1) = \{0\}$  and hence  $\Lambda_\lambda(\{0\}) = \{0\}$ . Therefore,  $A = \{0\}$  is a  $\Lambda_\lambda$ -set but it is not  $\lambda$ -open.

**Definition 3.5.** A subset  $A$  of a topological space  $X$  is said to be  $\Lambda_\lambda$ -closed (resp.  $\lambda$ -closed [1]) if  $A = L \cap F$ , where  $L$  is a  $\Lambda_\lambda$ -set (resp.  $\Lambda$ -set) and  $F$  is a closed set.

**Lemma 3.6.** For a subset of a topological space  $X$ , the following properties hold:

- (1) Every  $\lambda$ -locally closed set is  $\Lambda_\lambda$ -closed,
- (2) Every  $\lambda$ -closed set is  $\Lambda_\lambda$ -closed.

*Proof.* (1) By Lemma 3.3, every  $\lambda$ -open set is a  $\Lambda_\lambda$ -set and (1) holds.

(2) Let  $U$  be a  $\Lambda$ -set. Then,

$$U = \cap\{V : U \subseteq V, V \text{ is open}\} \supseteq \cap\{V : U \subset V, V \text{ is } \lambda\text{-open}\} \supseteq U$$

and hence  $U$  is a  $\Lambda_\lambda$ -set. Therefore, (2) holds.  $\square$

*Remark 3.7.* By Lemma 3.6, we obtain the following diagram.

DIAGRAM I

$$\begin{array}{ccc} \text{locally closed} & \Rightarrow & \lambda\text{-locally closed} \\ \Downarrow & & \Downarrow \\ \lambda\text{-closed} & \Rightarrow & \Lambda_\lambda\text{-closed} \end{array}$$

**Theorem 3.8.** *For a subset  $A$  of a topological space  $X$ , the following are equivalent:*

- (1)  $A$  is  $\Lambda_\lambda$ -closed;
- (2)  $A = U \cap \text{Cl}(A)$  for some  $\Lambda_\lambda$ -set  $U$ ;
- (3)  $A = \Lambda_\lambda(A) \cap \text{Cl}(A)$ .

*Proof.* (1)  $\Rightarrow$  (2): Let  $A$  be a  $\Lambda_\lambda$ -closed set. Then  $A = U \cap F$ , where  $U$  is a  $\Lambda_\lambda$ -set and  $F$  is a closed set. Thus, we have  $A \subseteq U \cap \text{Cl}(A) \subseteq U \cap \text{Cl}(F) = U \cap F = A$ . Therefore,  $A = U \cap \text{Cl}(A)$ .

(2)  $\Rightarrow$  (3): Let  $A = U \cap \text{Cl}(A)$  for some  $\Lambda_\lambda$ -set  $U$ . Since  $A \subseteq U$ , by Lemma 3.2  $\Lambda_\lambda(A) \subseteq \Lambda_\lambda(U) = U$  and hence  $A \subseteq \Lambda_\lambda(A) \cap \text{Cl}(A) \subseteq U \cap \text{Cl}(A) = A$ . Therefore, we obtain  $A = \Lambda_\lambda(A) \cap \text{Cl}(A)$ .

(3)  $\Rightarrow$  (1): Let  $A = \Lambda_\lambda(A) \cap \text{Cl}(A)$ . By Lemma 3.3,  $\Lambda_\lambda(A)$  is a  $\Lambda_\lambda$ -set and  $\text{Cl}(A)$  is closed. Therefore,  $A$  is  $\Lambda_\lambda$ -closed.  $\square$

**Definition 3.9.** Let  $X$  be a topological space. A subset  $A$  of  $X$  is said to be  $\lambda g$ -closed (resp.  $g$ -closed [8]) if  $\text{Cl}(A) \subseteq U$  whenever  $A \subseteq U$  and  $U$  is a  $\lambda$ -open (resp. open) set.

**Theorem 3.10.** *For a subset  $A$  of a topological space  $X$ , the following are equivalent:*

- (1)  $A$  is closed;
- (2)  $A$  is  $\lambda$ -locally closed and  $\lambda g$ -closed;
- (3)  $A$  is  $\Lambda_\lambda$ -closed and  $\lambda g$ -closed.

*Proof.* (1)  $\Rightarrow$  (2): Let  $A$  be closed in  $X$ . Since  $A = X \cap A$  and  $X$  is a  $\Lambda_\lambda$ -set,  $A$  is  $\lambda$ -locally closed. Let  $U$  be any  $\lambda$ -open set containing  $A$ . Then  $\text{Cl}(A) = A \subseteq U$  and hence  $A$  is  $\lambda g$ -closed.

(2)  $\Rightarrow$  (3): By Lemma 3.6, every  $\lambda$ -locally closed set is  $\Lambda_\lambda$ -closed.

(3)  $\Rightarrow$  (1): Let  $A$  be  $\Lambda_\lambda$ -closed and  $\lambda g$ -closed. Since  $A$  is  $\Lambda_\lambda$ -closed,  $A = P \cap L$ , where  $P$  is a  $\Lambda_\lambda$ -set and  $L$  is closed in  $X$ . Let  $V$  be any  $\lambda$ -open set containing  $A$ . Since  $A$  is  $\lambda g$ -closed,  $\text{Cl}(A) \subseteq V$  and hence  $\text{Cl}(A) \subseteq \bigcap \{V : A \subseteq V, V \text{ is } \lambda\text{-open}\} = \Lambda_\lambda(A)$ . Therefore,  $\text{Cl}(A) \subseteq \Lambda_\lambda(A) \subseteq \Lambda_\lambda(P) = P$ . On the other hand,  $A \subseteq L$  and  $\text{Cl}(A) \subseteq \text{Cl}(L) = L$ . Therefore, we obtain  $\text{Cl}(A) \subseteq P \cap L = A$ . Thus  $A$  is closed.  $\square$

**Theorem 3.11.** *Let  $X$  be a topological space. If  $A_\alpha$  is a  $\Lambda_\lambda$ -closed set for each  $\alpha \in \Delta$ , then  $\bigcap_{\alpha \in \Delta} A_\alpha$  is  $\Lambda_\lambda$ -closed.*

*Proof.* Let  $A_\alpha$  be a  $\Lambda_\lambda$ -closed set for each  $\alpha \in \Delta$ . Then  $A_\alpha = U_\alpha \cap F_\alpha$ , where  $U_\alpha$  is a  $\Lambda_\lambda$ -set and  $F_\alpha$  is a closed set for each  $\alpha \in \Delta$ . By Lemma 3.3,  $\bigcap_{\alpha \in \Delta} U_\alpha$  is a  $\Lambda_\lambda$ -set,  $\bigcap_{\alpha \in \Delta} F_\alpha$  is closed and  $\bigcap_{\alpha \in \Delta} A_\alpha = (\bigcap_{\alpha \in \Delta} U_\alpha) \cap (\bigcap_{\alpha \in \Delta} F_\alpha)$ . Therefore,  $\bigcap_{\alpha \in \Delta} A_\alpha$  is  $\Lambda_\lambda$ -closed.  $\square$

## 4. DECOMPOSITIONS OF CONTINUITY

In this section, we obtain the decompositions of continuity.

**Definition 4.1.** A function  $f : X \rightarrow Y$  is said to be

- (1)  $\lambda$ -*LC-continuous* if  $f^{-1}(V)$  is  $\lambda$ -locally closed in  $X$  for any closed set  $V$  of  $Y$ ,
- (2)  $\Lambda_\lambda$ -*continuous* if  $f^{-1}(V)$  is  $\Lambda_\lambda$ -closed in  $X$  for any closed set  $V$  of  $Y$ ,
- (3)  $\lambda g$ -*continuous* if  $f^{-1}(V)$  is  $\lambda g$ -closed in  $X$  for any closed set  $V$  of  $Y$ .

**Theorem 4.2.** For a function  $f : X \rightarrow Y$ , the following are equivalent:

- (1)  $f$  is continuous;
- (2)  $f$  is  $\lambda$ -*LC-continuous* and  $\lambda g$ -*continuous*;
- (3)  $f$  is  $\Lambda_\lambda$ -*continuous* and  $\lambda g$ -*continuous*.

*Proof.* This is an immediate consequence of Theorem 3.10 □

**Remark 4.3.** The following facts are shown by Examples 4.4 and 4.5 and Remark 4.6:

- (1)  $\lambda$ -*LC-continuity* and  $\lambda g$ -*continuity* are independent of each other,
- (2)  $\Lambda_\lambda$ -*continuity* and  $\lambda g$ -*continuity* are independent of each other.

**EXAMPLE 4.4.** Let  $X = Y = \{a, b, c, d\}$ ,  $\tau = \sigma = \{\emptyset, X, \{a\}\}$ . Then  $C(X) = C(Y) = \{\emptyset, \{b, c, d\}\}$  and  $\lambda$ -open sets in  $X$  (resp.  $Y$ ) are:  $\emptyset, X, \{a\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, b, c\}, \{a, c, d\}, \{a, b, d\}$ .  $\lambda$ -locally closed sets in  $X$  (resp.  $Y$ ) are:  $\emptyset, X, \{a\}, \{a, b\}, \{a, c\}, \{a, d\}, \{a, b, c\}, \{a, c, d\}, \{a, b, d\}, \{b, c, d\}, \{b, c\}, \{c, d\}, \{b, d\}, \{b\}, \{c\}, \{d\}$ . Define a function  $f : X \rightarrow Y$  by  $f(a) = c, f(b) = b, f(c) = d, f(d) = a$ . Then we have the following:

- (1) Since  $f^{-1}(\{b, c, d\}) = \{a, b, c\}$ , then  $f$  is not continuous.
- (2) Since  $f^{-1}(\{b, c, d\}) = \{a, b, c\}$ , then  $f$  is  $\lambda$ -*LC-continuous*.
- (3) Since  $Cl(\{a, b, c\}) = X$  (i.e.  $\{a, b, c\}$  is not  $\lambda g$ -closed), then  $f$  is not  $\lambda g$ -continuous.
- (4) Since  $\{a, b, c\} \subseteq \cap\{U : \{a, b, c\} \subseteq U, U \text{ is } \lambda\text{-open}\} = \{a, b, c\}$  and  $\{a, b, c\} = \{a, b, c\} \cap X = \{a, b, c\}$ , then  $\{a, b, c\}$  is  $\Lambda_\lambda$ -closed. Thus  $f$  is  $\Lambda_\lambda$ -continuous.

**EXAMPLE 4.5.** Let  $X = Y = \{a, b, c, d\}$ ,  $\tau = \sigma = \{\emptyset, X, \{a, b\}, \{c, d\}\}$ . Then  $C(X) = C(Y) = \{\emptyset, X, \{a, b\}, \{c, d\}\}$  and  $\lambda$ -open sets in  $X$  (resp.  $Y$ ) are:  $\emptyset, X, \{a, b\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}$ . And  $\lambda$ -locally closed sets in  $X$  (resp.  $Y$ ) are:  $\emptyset, X, \{a, b\}, \{c, d\}, \{a, b, c\}, \{a, b, d\}, \{a, c, d\}, \{b, c, d\}, \{a\}, \{b\}, \{c\}, \{d\}$ . Define  $g : X \rightarrow Y$  by  $g(a) = c, g(b) = b, g(c) = a, g(d) = d$ . Then we have the following:

- (1) Since  $g^{-1}(\{c, d\}) = \{a, d\}$ , then  $g$  is not a continuous function.
- (2) Since  $g^{-1}(\{c, d\}) = \{a, d\}$ , it is not a  $\lambda$ -locally closed set in  $X$ . Then  $g$  is not a  $\lambda$ -*LC-continuous* function.
- (3) Since  $g^{-1}(\{a, b\}) = \{b, c\} \subseteq \cap\{U : \{b, c\} \subseteq U, U \text{ is } \lambda\text{-open in } X\} =$

$\{b, c\} \cap X = \{b, c\}$  and  $g^{-1}(\{c, d\}) = \{a, d\} = \cap \{U : \{a, d\} \subseteq U, U \text{ is } \lambda\text{-open in } X\} = \{a, d\} \cap X = \{a, d\}$  are  $\Lambda_\lambda$ -closed, then  $\Lambda_\lambda$ -continuous.

*Remark 4.6.* (1) If every  $\lambda g$ -continuous function is  $\lambda$ -LC-continuous, then it is continuous from Theorem 4.2 This is not true from Example 4.4(1).

(2) If every  $\lambda g$ -continuous function is  $\Lambda_\lambda$ -continuous, then it is continuous from Theorem 4.2. This not true from Example 4.5(1).

#### ACKNOWLEDGMENTS

The authors wish to thank the referees for their valuable comments.

#### REFERENCES

1. F. G. Arenas, J. Dontchev, M. Ganster, On  $\lambda$ -Sets and the Dual of Generalized Continuity, *Questions Answers General Topology*, **15**, (1997), 3–13.
2. R. A. Borzooei, G. R. Rezaei, N. Kouhestani, On (semi) Topological  $BL$ -Algebras, *Iranian Journal of Mathematical Sciences and Informatics*, **6**(1), (2011), 59–77.
3. N. Bourbaki, *General Topology*, Chapters 1-4, Springer-Verlag, 1989.
4. J. Dontchev, On Superconnected Spaces, *Serdica-Bulgaricae Mathematicae Publicationes*, **20**, (1994), 345–350.
5. A. A. Estaji,  $z$ -Weak Ideals and Prime Weak Ideals, *Iranian Journal of Mathematical Sciences and Informatics*, **7**(2), (2012), 53–62.
6. M. Ganster, I. L. Reilly, Locally Closed Sets and  $LC$ -continuous Functions, *Internat. J. Math. Math. Sci.*, **12**(3), (1989), 417–424.
7. N. Levine, Semi-open Sets and Semi-continuity in Topological Spaces, *Amer. Math. Monthly*, **70**, (1963), 36–41.
8. N. Levine, Generalized Closed Sets in Topology, *Rend. Circ. Mat. Palermo (2)*, **19**, (1970), 89–96.
9. H. Maki, Generalized  $\Lambda$ -Sets and the Associated Closure Operator, *The Special Issue in Commemoration of Prof. Kazusada IKEDA's Retirement*, (1986), 139–146.
10. A. S. Mashhour, M. E. Abd El-Monsef, S. N. El-Deeb, On Precontinuous and Weak Precontinuous Mappings, *Proc. Math. Phys. Soc. Egypt.*, **53**, (1982), 47–53.
11. O. Njåstad, On Some Classes of Nearly Open Sets, *Pacific J. Math.*, **15**, (1965), 961–970.
12. T. Roudari, L. Torkzadeh, A Topology on  $BCK$ -Algebra via Left and Right Stabilizers, *Iranian Journal of Mathematical Sciences and Informatics*, **4**(2), (2009), 1–8.